

**This Page Is Inserted by IFW Operations  
and is not a part of the Official Record**

## **BEST AVAILABLE IMAGES**

**Defective images within this document are accurate representations of the original documents submitted by the applicant.**

**Defects in the images may include (but are not limited to):**


- **BLACK BORDERS**
- **TEXT CUT OFF AT TOP, BOTTOM OR SIDES**
- **FADED TEXT**
- **ILLEGIBLE TEXT**
- **SKEWED/SLANTED IMAGES**
- **COLORED PHOTOS**
- **BLACK OR VERY BLACK AND WHITE DARK PHOTOS**
- **GRAY SCALE DOCUMENTS**

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning documents *will not* correct images,  
please do not report the images to the  
Image Problem Mailbox.**



## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<p>(51) International Patent Classification 7 : C07D 217/04, 471/04, A61K 31/472, 31/4725, 31/4375, C07D 405/12, 409/12, 401/12, 217/06, 217/02, 413/12, 401/06 // (C07D 471/04, 221:00, 221:00)</p>	<p>A1</p>	<p>(11) International Publication Number: <b>WO 00/07993</b>  (43) International Publication Date: 17 February 2000 (17.02.00)</p>
<p>(21) International Application Number: PCT/EP99/05583 (22) International Filing Date: 3 August 1999 (03.08.99)  (30) Priority Data: 9816984.0 5 August 1998 (05.08.98) GB  (71) Applicant (for all designated States except US): SMITHKLINE BEECHAM PLC [GB/GB]; New Horizons Court, Brentford, Middlesex TW8 9EP (GB).  (72) Inventors; and (75) Inventors/Applicants (for US only): COULTON, Steven [GB/GB]; SmithKline Beecham Pharmaceuticals, New Frontiers Science Park South, Third Avenue, Harlow, Essex CM19 5AW (GB). HARLING, John, David [GB/GB]; SmithKline Beecham Pharmaceuticals, New Frontiers Science Park South, Third Avenue, Harlow, Essex CM19 5AW (GB). PORTER, Roderick, Alan [GB/GB]; SmithK- line Beecham Pharmaceuticals, New Frontiers Science Park South, Third Avenue, Harlow, Essex CM19 5AW (GB). THOMPSON, Mervyn [GB/GB]; SmithKline Beecham Pharmaceuticals, New Frontiers Science Park South, Third Avenue, Harlow, Essex CM19 5AW (GB).</p>	<p>(74) Agent: RUSSELL, Brian, John; SmithKline Beecham, Corpo- rate Intellectual Property, Two New Horizons Court, Brent- ford, Middlesex TW8 9EP (GB).  (81) Designated States: CA, JP, US, European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).  <b>Published</b> <i>With international search report.</i></p>	
<p>(54) Title: SUBSTITUTED ISOQUINOLEINES AND THEIR USE AS ANTICONVULSIVANTS</p>		
<div style="text-align: center;">  <p>(I)</p> </div>		
<p>(57) Abstract</p> <p>Compounds of formula (I) including tetrahydroisoquinolinyl cinnamides and acrylamides are indicated to be useful for the treatment and/or prevention of anxiety, mania, depression, panic disorders and/or aggression, disorders associated with a subarachnoid haemorrhage or neural shock, the effects associated with withdrawal from substances of abuse such as cocaine, nicotine, alcohol and benzodiazepines, disorders treatable and/or preventable with anti-convulsive agents, such as epilepsy including post-traumatic epilepsy, Parkinson's disease, psychosis, migraine, cerebral ischaemia, Alzheimer's disease and other degenerative diseases such as Huntingdon's chorea, schizophrenia, obsessive compulsive disorders (OCD), neurological deficits associated with AIDS, sleep disorders (including circadian rhythm disorders, insomnia and narcolepsy), tics (e.g. Gilles de la Tourette's syndrome), traumatic brain injury, tinnitus, neuralgia, especially trigeminal neuralgia, neuropathic pain, dental pain, cancer pain, inappropriate neuronal activity resulting in neurodysesthesias in diseases such as diabetes, multiple sclerosis (MS) and motor neurone disease, ataxias, muscular rigidity (spasticity), temporomandibular joint dysfunction, and amyotrophic lateral sclerosis (ALS).</p>		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						


## SUBSTITUTED ISOQUINOLEINES AND THEIR USE AS ANTICONVULSIVANTS

This invention relates to novel compounds, to processes for preparing them, and to their use as therapeutic agents.

- 5 It has now been surprisingly found that cinnamide and acrylamide compounds of formula (I) below possess anti-convulsant activity and are therefore believed to be useful in the treatment and/or prevention of anxiety, mania, depression, panic disorders and/or aggression, disorders associated with a subarachnoid haemorrhage or neural shock, the effects associated with withdrawal  
10 from substances of abuse such as cocaine, nicotine, alcohol and benzodiazepines, disorders treatable and/or preventable with anti-convulsive agents, such as epilepsy including post-traumatic epilepsy, Parkinson's disease, psychosis, migraine, cerebral ischaemia, Alzheimer's disease and other degenerative diseases such as Huntingdon's chorea, schizophrenia, obsessive compulsive disorders  
15 (OCD), neurological deficits associated with AIDS, sleep disorders (including circadian rhythm disorders, insomnia & narcolepsy), tics (e.g. Gilles de la Tourette's syndrome), traumatic brain injury, tinnitus, neuralgia, especially trigeminal neuralgia, neuropathic pain, dental pain, cancer pain, inappropriate neuronal activity resulting in neurodysthesis in diseases such as diabetes, multiple sclerosis (MS) and motor neurone disease, ataxias, muscular rigidity  
20 (spasticity), temporomandibular joint dysfunction, and amyotrophic lateral sclerosis (ALS).

Accordingly, the present invention provides a compound of formula (I) or pharmaceutically acceptable salt thereof:

25



in which

- 30 Z is a carbocyclic or heterocyclic or a fused carbocyclic or heterocyclic ring containing at least one aromatic ring;

X is CH or N;

Y is hydrogen, C<sub>1-6</sub>alkyl, or a halogen;

P is -CH=CH- and Q is -NR<sup>1</sup>-, or;

P is -CH=CH- and Q is -NR<sup>1</sup>CH<sub>2</sub>-, or;

- P is -NH- and Q is -CR<sup>1a</sup>=CH-;  
 R<sup>1</sup> is hydrogen, phenylC<sub>1-6</sub> alkyl, or C<sub>1-6</sub> alkyl;  
 R<sup>1a</sup> is hydrogen, halogen, phenylC<sub>1-6</sub> alkyl, or C<sub>1-6</sub> alkyl;  
 R<sup>2</sup> is hydrogen or up to three substituents selected from halogen, NO<sub>2</sub>, CN, N<sub>3</sub>,  
 5 CF<sub>3</sub>O-, CF<sub>3</sub>S-, CF<sub>3</sub>CO-, CF<sub>3</sub>SO<sub>2</sub>, C<sub>1-6</sub>alkyl,  
 C<sub>1-6</sub>alkenyl, C<sub>1-6</sub>alkynyl, C<sub>1-6</sub>perfluoroalkyl, C<sub>3-6</sub>cycloalkyl,  
 C<sub>3-6</sub>cycloalkyl-C<sub>1-4</sub>alkyl-, C<sub>1-6</sub>alkylO-, C<sub>1-6</sub>alkylCO-, C<sub>3-6</sub>cycloalkylO-,  
 C<sub>3-6</sub>cycloalkylCO-, C<sub>3-6</sub>cycloalkyl-C<sub>1-4</sub>alkylO-, C<sub>3-6</sub>cycloalkyl-C<sub>1-4</sub>alkylCO-,  
 phenyl, phenoxy, benzyloxy, benzoyl, phenyl-C<sub>1-4</sub>alkyl-, C<sub>1-6</sub>alkylS-,  
 10 C<sub>1-6</sub>alkylSO<sub>2</sub>-, or 1,3-oxazol-5-yl, (C<sub>1-4</sub>alkyl)<sub>2</sub>NSO<sub>2</sub>-, (C<sub>1-4</sub>alkyl)NHSO<sub>2</sub>-,  
 (C<sub>1-4</sub>alkyl)<sub>2</sub>NCO-, (C<sub>1-4</sub>alkyl)NHCO- or CONR<sup>4</sup>R<sup>5</sup>, CO<sub>2</sub>R<sup>4</sup>;  
 or -NR<sup>4</sup>R<sup>6</sup> or NHCOR<sup>4</sup>  
 where R<sup>4</sup> and R<sup>5</sup> are each independently hydrogen or C<sub>1-4</sub> alkyl, and;  
 R<sup>6</sup> is hydrogen, C<sub>1-4</sub>alkyl, formyl, -CO<sub>2</sub>C<sub>1-4</sub>alkyl, or -COC<sub>1-4</sub>alkyl;  
 15 or two R<sup>2</sup> groups are linked together to form a carbocyclic ring that is saturated or  
 unsaturated and unsubstituted or substituted by -OH or =O or a heterocyclic ring  
 that is saturated or unsaturated;  
 or when P is -CH=CH- and Q is -NR<sup>1</sup>CH<sub>2</sub>-, R<sup>1</sup> and an R<sup>2</sup> are linked together to  
 form a saturated or unsaturated carbocyclic or heterocyclic ring;  
 20 or when P is -CH=CH- and Q is -NR<sup>1</sup>-, R<sup>1</sup> and an R<sup>2</sup> are linked together to form a  
 saturated or unsaturated carbocyclic or heterocyclic ring, and;  
 R<sup>3</sup> is hydrogen, phenylC<sub>1-6</sub> alkyl, C<sub>1-6</sub> alkyl, C<sub>1-6</sub> alkylOCO-, C<sub>1-6</sub>alkylCO-,  
 formyl, CF<sub>3</sub>CO- or C<sub>1-6</sub>alkylSO<sub>2</sub>-, hydroxyC<sub>1-6</sub>alkyl, or C<sub>1-6</sub>alkoxyC<sub>1-6</sub>alkyl.  
 R<sup>7</sup> is hydrogen or C<sub>1-6</sub> alkyl;  
 25 R<sup>8</sup> is hydrogen or C<sub>1-6</sub> alkyl;  
 R<sup>9</sup> is hydrogen or C<sub>1-6</sub> alkyl;  
 R<sup>10</sup> is hydrogen or C<sub>1-6</sub> alkyl;  
 R<sup>11</sup> is hydrogen or C<sub>1-6</sub> alkyl, and;  
 R<sup>12</sup> is hydrogen or C<sub>1-6</sub> alkyl.  
 30 In the formula (I), alkyl groups, including alkyl groups that are part of  
 another moiety, may be straight chain or branched. Aromatic rings, especially  
 phenyl groups, including rings that are part of another moiety, may optionally be  
 substituted with one or more independently selected halogen, C<sub>1-6</sub> alkyl, C<sub>1-6</sub>

alkoxy or C<sub>1-6</sub> alkylcarbonyl groups. Suitable halo substituents include fluoro, chloro, iodo and bromo. Suitable C<sub>3-6</sub> cycloalkyl groups include cyclopropyl, cyclobutyl, cyclopentyl, and cyclohexyl groups.

When used herein the terms "heterocyclyl" and "heterocyclic" suitably include, unless otherwise defined, aromatic and non-aromatic, single and fused, rings suitably containing up to four heteroatoms in each ring, each of which is selected from oxygen, nitrogen and sulphur, which rings may be unsubstituted or substituted by, for example, up to three substituents. Each heterocyclic ring suitably has from 4 to 7, preferably 5 or 6, ring atoms. A fused heterocyclic ring system may include carbocyclic rings and need include only one heterocyclic ring.

When ring Z is heterocyclic, Z may be for example furanyl, thiophenyl, indolinyl or indazolinyl. Preferably Z is phenyl.

Linked R<sup>2</sup> groups and linked R<sup>1</sup> and R<sup>2</sup> groups are typically such as to form a 5 or 6 membered ring fused to the ring to which the R<sup>2</sup> groups are appended. Thus when Z is phenyl, the linked R<sup>2</sup> groups or linked R<sup>1</sup> and R<sup>2</sup> groups may create fused rings such that the moiety Q is tetrahydroquinolinyl, tetrahydroisoquinolinyl or dihydroindolinyl.

Preferably a substituent for a heterocyclyl group is selected from halogen, (C<sub>1-6</sub>)alkyl, aryl(C<sub>1-6</sub>)alkyl, (C<sub>1-6</sub>)alkoxy, (C<sub>1-6</sub>)alkoxy(C<sub>1-6</sub>)alkyl, halo(C<sub>1-6</sub>)alkyl, hydroxy, amino, mono- and di-N-(C<sub>1-6</sub>)alkyl-amino, acylamino, carboxy, carboxy salts, carboxy esters, carbamoyl, mono- and di-N-(C<sub>1-6</sub>)alkylcarbonyl, aryloxy, carbonyl, (C<sub>1-6</sub>)alkoxy, carbonyl(C<sub>1-6</sub>)alkyl, aryloxy groups, ureido, guanidino, sulphonylamino, aminosulphonyl, (C<sub>1-6</sub>)alkylthio, (C<sub>1-6</sub>)alkylsulphinyl, (C<sub>1-6</sub>)alkylsulphonyl, heterocyclyl and heterocyclyl(C<sub>1-6</sub>)alkyl.

It should be appreciated that the compounds of formula (I) may have chiral carbon atoms and therefore may exist as enantiomers. The present invention extends to each enantiomer and to mixtures thereof including racemates.

Preferably where P is -CH=CH- or Q is CR<sup>1a</sup>=CH the compound exists as the E isomer.

A suitable group of compounds of formula (I) have:

R<sup>1</sup> as hydrogen, fluoro, methyl, ethyl or propyl;  
 R<sup>2</sup> as hydrogen or one or more of methyl, ethyl, *n*-butyl, phenyl, *iso*-propyl, *t*-butyl, methoxy, ethoxy, *n*-propoxy, *iso*-propoxy, *n*-butoxy, phenoxy, benzyloxy, bromo, chloro, iodo, fluoro, nitro, cyano, acetyl, pivaloyl, *iso*-butyryl, benzoyl, trifluoromethyl, trifluoromethoxy, trifluoroacetyl, amino, acetylamino, methylthio, oxazolo, methylsulfonyl, *n*-propylsulfonyl, isopropylsulfonyl or dimethylsulfamoyl;

R<sup>3</sup> as hydrogen, methyl, ethyl, propyl, benzyl, *t*-butyloxycarbonyl or trifluoroacetyl.

Suitable linked R<sup>2</sup> groups include -CH=CH-NH-.

- 5     Suitable linked R<sup>1</sup> and R<sup>2</sup> groups are ethylene, propylene, 1,1-dimethylethylene when Q is -NR<sup>1</sup>; or suitable linked R<sup>1</sup> and R<sup>2</sup> groups are ethylene, propylene, 1,1-dimethylethylene when Q is -NR<sup>1</sup>CH<sub>2</sub>.

In a particular group of compounds of formula (I),

R<sup>1</sup> is hydrogen, fluoro or methyl;

- 10     R<sup>2</sup> is hydrogen or one or more of methyl, ethyl, *t*-butyl, methoxy, methoxycarbonyl, methylcarbonyl, ethylcarbonyl, methylamido, acetylamino, methylsulfonyl, oxazole, trifluoromethyl, cyano, chloro, fluoro, or nitro;

R<sup>3</sup> is hydrogen, methyl, ethyl, *n*-propyl, benzyl or *t*-butyloxycarbonyl.

Examples of compounds of one aspect of formula (I) are:

- 15     E-N-(2-Methyl-1,2,3,4-tetrahydroisoquinolin-7-yl)-3-nitrocinnamide;  
 E-N-(2-Methyl-1,2,3,4-tetrahydroisoquinolin-7-yl)-3-trifluoromethylcinnamide;  
 E-N-(2-Methyl-1,2,3,4-tetrahydroisoquinolin-7-yl)-3-cinnamide hydrochloride;  
 E-N-(2-Methyl-1,2,3,4-tetrahydroisoquinolin-7-yl)-2-methoxycinnamide;  
 E-N-(2-Methyl-1,2,3,4-tetrahydroisoquinolin-7-yl)-4-chlorocinnamide;  
 E-N-(2-Methyl-1,2,3,4-tetrahydroisoquinolin-7-yl)-3-chlorocinnamide;  
 20     E-N-(2-Methyl-1,2,3,4-tetrahydroisoquinolin-7-yl)-3-methoxycinnamide;  
 E-N-(2-Methyl-1,2,3,4-tetrahydroisoquinolin-7-yl)- $\alpha$ -methylcinnamide;  
 E-N-(2-Methyl-1,2,3,4-tetrahydroisoquinolin-7-yl)-2-chlorocinnamide;  
 E-N-(2-Methyl-1,2,3,4-tetrahydroisoquinolin-7-yl)-4-methoxycinnamide;  
 E-N-(6-methyl-5,6,7,8-tetrahydro[1,6]naphthyridin-3-yl)-3-phenylacrylamide;  
 25     E-3-Furan-2-yl-N-(2-methyl-1,2,3,4-tetrahydroisoquinolin-7-yl)acrylamide;  
 E-N-(2-methyl-1,2,3,4-tetrahydroisoquinolin-7-yl)-3-thiophen-2-ylacrylamide;  
 E-N-(2-Methyl-1,2,3,4-tetrahydroisoquinolin-7-yl)-2, 4-dichlorocinnamide;  
 Z-N-(2-Methyl-1,2,3,4-tetrahydroisoquinolin-7-yl)-2-methoxycinnamide;  
 E-3-Indolin-5-yl-N-(2-Methyl-1,2,3,4-tetrahydroisoquinolin-7-yl)-acrylamide;  
 30     E-3-(1-Methyl-Indolin-2-yl)-N-(2-Methyl-1,2,3,4-tetrahydroisoquinolin-7-yl)-acrylamide;  
 E-N-(2-Methyl-1,2,3,4-tetrahydroisoquinolin-7-yl)-3-chloro-4-methoxycinnamide;  
 E-N-(2-Methyl-1,2,3,4-tetrahydroisoquinolin-7-yl)-4-methylsulphonylcinnamide;  
 E-N-methyl-3-[2-(2-methyl-1,2,3,4-tetrahydroisoquinolin-7-ylcarbonyl)vinyl]  
 35     benzamide;  
 E-3-(Indazolin-3-yl)-N-(2-Methyl-1,2,3,4-tetrahydroisoquinolin-7-yl)-acrylamide;  
 E-N-(2-Methyl-1,2,3,4-tetrahydroisoquinolin-7-yl)-2-methylcinnamide;  
 E-N-(2-Methyl-1,2,3,4-tetrahydroisoquinolin-7-yl)-2-nitrocinnamide;  
 E-N-(2-Methyl-1,2,3,4-tetrahydroisoquinolin-7-yl)-2-trifluoromethylcinnamide;

- E-N-(2-Methyl-1,2,3,4-tetrahydroisoquinolin-7-yl)-2-ethoxycinnamide;  
E-N-(2-Methyl-1,2,3,4-tetrahydroisoquinolin-7-yl)-2-chloro-4-fluorocinnamide;  
E-N-(2-Methyl-1,2,3,4-tetrahydroisoquinolin-7-yl)-2-chloro-6-fluorocinnamide;  
E-N-(2-Methyl-1,2,3,4-tetrahydroisoquinolin-5-yl)-4-chlorocinnamide;  
5 E-N-(2-Methyl-1,2,3,4-tetrahydroisoquinolin-5-yl)-cinnamide;  
E-N-(2-Methyl-1,2,3,4-tetrahydroisoquinolin-5-yl)-3-chlorocinnamide;  
E-N-(2-Methyl-1,2,3,4-tetrahydroisoquinolin-5-yl)-2-chlorocinnamide;  
E-N-(2-Methyl-1,2,3,4-tetrahydroisoquinolin-5-yl)-3-acetylcinnamide;  
E-N-(2-Methyl-1,2,3,4-tetrahydroisoquinolin-5-yl)-2-bromocinnamide;  
10 E-N-(2-Methyl-1,2,3,4-tetrahydroisoquinolin-5-yl)-2-methylcinnamide;  
E-N-(2-Methyl-1,2,3,4-tetrahydroisoquinolin-5-yl)-4-ethoxycinnamide;  
E-N-(2-Methyl-1,2,3,4-tetrahydroisoquinolin-5-yl)-2-methoxycinnamide;  
E-N-(2-Methyl-1,2,3,4-tetrahydroisoquinolin-5-yl)-5-bromo-2-methoxycinnamide;  
E-2-Cyano-N-(2-Methyl-1,2,3,4-tetrahydroisoquinolin-5-yl)cinnamide;  
15 N-(8-Chloro-2-trifluoroacetyl-1,2,3,4-tetrahydroisoquinolin-7-yl)-2-chlorocinnamide;  
N-(8-Chloro-1,2,3,4-tetrahydroisoquinolin-7-yl)-2-chlorocinnamide;  
N-(8-Chloro-2-methyl-1,2,3,4-tetrahydroisoquinolin-7-yl)-2-chlorocinnamide;  
N-(8-Bromo-2-trifluoroacetyl-1,2,3,4-tetrahydroisoquinolin-7-yl)-2-chlorocinnamide;  
20 E-N-(2-Methyl-1,2,3,4-tetrahydroisoquinolin-7-yl)- $\alpha$ -fluorocinnamide;  
E-N-(8-Bromo-1,2,3,4-tetrahydroisoquinolin-7-yl)-2-chlorocinnamide;  
E-N-(8-Bromo-2-methyl-1,2,3,4-tetrahydroisoquinolin-7-yl)-2-chlorocinnamide;  
E-N-(2,4,4-Trimethyl-1,2,3,4-tetrahydroisoquinolin-7-yl)-2-chlorocinnamide;  
25 E-N-(8-Ethyl-2,4,4-trimethyl-1,2,3,4-tetrahydroisoquinolin-7-yl)-2-methylcinnamide;  
E-N-(8-Ethyl-2,4,4-trimethyl-1,2,3,4-tetrahydroisoquinolin-7-yl)-2-chloro-6-fluorocinnamide;  
E-N-(8-Ethyl-2,4,4-trimethyl-1,2,3,4-tetrahydroisoquinolin-7-yl)-2-trifluoromethylcinnamide;  
30 E-N-(8-Ethyl-2,4,4-trimethyl-1,2,3,4-tetrahydroisoquinolin-7-yl)-2-chloro-4-fluorocinnamide;  
E-N-(8-Ethyl-2,4,4-trimethyl-1,2,3,4-tetrahydroisoquinolin-7-yl)-2-chlorocinnamide;  
35 E-N-(1,1,2-Trimethyl-1,2,3,4-tetrahydroisoquinolin-7-yl)-2-chlorocinnamide;  
E-N-(1,2,4,4-Tetramethyl-1,2,3,4-tetrahydroisoquinolin-7-yl)-2-chlorocinnamide;  
E-N-(8-Ethyl-2,4,4-trimethyl-1,2,3,4-tetrahydroisoquinolin-7-yl)-2-ethoxycinnamide;



- E-N-(8-Chloro-2,4,4-trimethyl-1,2,3,4-tetrahydroisoquinolin-7-yl)-2-chlorocinnamide;  
 E-N-(8-Ethyl-2,4,4-trimethyl-1,2,3,4-tetrahydroisoquinolin-7-yl)-3-chloro-4-methoxycinnamide;
- 5 E-N-(8-Ethyl-2,4,4-trimethyl-1,2,3,4-tetrahydroisoquinolin-7-yl)-2-cyanocinnamide;  
 E-N-(8-Methyl-2,4,4-trimethyl-1,2,3,4-tetrahydroisoquinolin-7-yl)-2-chlorocinnamide;  
 E-N-(8-Ethyl-2,4,4-trimethyl-1,2,3,4-tetrahydroisoquinolin-7-yl)-2-
- 10 acetylcinnamide;  
 E-N-(8-Ethyl-4,4-dimethyl-1,2,3,4-tetrahydroisoquinolin-7-yl)-2-chlorocinnamide;  
 E-N-(8-Ethyl-4,4-dimethyl-1,2,3,4-tetrahydroisoquinolin-7-yl)-2-cyanocinnamide;  
 E-N-(8-Chloro-2,3,3-trimethyl-1,2,3,4-tetrahydroisoquinolin-7-yl)-2-chlorocinnamide;
- 15 E-N-(5-Bromo-1,2,3,4-tetrahydroisoquinolin-7-yl)-2-chlorocinnamide;  
 E-N-(5,6,7,8-Tetrahydro-6-methyl[1,6]naphthyridin-3-yl)-cinnamide, and;  
 E-N-(5,6,7,8-Tetrahydro-6,8,8-trimethyl[1,6]naphthyridin-3-yl)-2-chlorocinnamide.
- Examples of compounds of another aspect of formula (I) are:
- 20 E-N-(4-Methoxyphenyl)-3-(1,2,3,4-tetrahydroisoquinolin-7-yl)acrylamide;  
 E-N-Phenyl-3-(1,2,3,4-tetrahydroisoquinolin-7-yl)acrylamide;  
 E-N-Phenyl-3-(2-methyl-1,2,3,4-tetrahydroisoquinolin-7-yl)acrylamide;  
 E-N-Phenyl-3-(2-benzyl-1,2,3,4-tetrahydroisoquinolin-7-yl)acrylamide;  
 E-N-Phenyl-3-(2-n-propyl-1,2,3,4-tetrahydroisoquinolin-7-yl)acrylamide;
- 25 E-N-Phenyl-3-(2-ethyl-1,2,3,4-tetrahydroisoquinolin-7-yl)acrylamide;  
 E-N-(3-Cyanophenyl)-3-(1,2,3,4-tetrahydroisoquinolin-7-yl)acrylamide;  
 E-N-(3-Cyanophenyl)-3-(2-methyl-1,2,3,4-tetrahydroisoquinolin-7-yl)acrylamide;  
 E-N-(2-Chlorophenyl)-3-(2-methyl-1,2,3,4-tetrahydroisoquinolin-7-yl)acrylamide;  
 E-N-(2-Methoxyphenyl)-3-(2-methyl-1,2,3,4-tetrahydroisoquinolin-7-
- 30 yl)acrylamide;  
 E-N-(3-Methoxyphenyl)-3-(2-methyl-1,2,3,4-tetrahydroisoquinolin-7-yl)acrylamide;  
 E-N-(3-Chlorophenyl)-3-(2-methyl-1,2,3,4-tetrahydroisoquinolin-7-yl)acrylamide;  
 E-N-(4-Chlorophenyl)-3-(2-methyl-1,2,3,4-tetrahydroisoquinolin-7-yl)acrylamide;
- 35 E-N-Methyl-N-benzyl-3-(2-methyl-1,2,3,4-tetrahydroisoquinolin-7-yl)acrylamide;  
 E-N-(3-Nitrophenyl)-3-(2-methyl-1,2,3,4-tetrahydroisoquinolin-7-yl)acrylamide;  
 E-N-Methyl-N-phenyl-3-(2-methyl-1,2,3,4-tetrahydroisoquinolin-7-yl)acrylamide;  
 E-N-(3-Carbomethoxyphenyl)-3-(2-methyl-1,2,3,4-tetrahydroisoquinolin-7-yl)acrylamide;

- E-N-Methyl-3-[3-(2-methyl-1,2,3,4-tetrahydroisoquinolin-7-yl)acryloylamino]benzamide;  
 E-N-(3-N-Methylsulphonylphenyl)-3-(2-methyl-1,2,3,4-tetrahydroisoquinolin-7-yl)acrylamide;  
 5 E-N-(1,3-Oxazol-5-ylphenyl)-3-(2-methyl-1,2,3,4-tetrahydroisoquinolin-7-yl)acrylamide;  
 E-N-(3-Acetylaminophenyl)-3-(2-methyl-1,2,3,4-tetrahydroisoquinolin-7-yl)acrylamide;  
 E-N-(3-Ethylphenyl)-3-(2-methyl-1,2,3,4-tetrahydroisoquinolin-7-yl)acrylamide;  
 10 E-N-(3-Methylphenyl)-3-(2-methyl-1,2,3,4-tetrahydroisoquinolin-7-yl)acrylamide;  
 E-N-(3-tert-Butylphenyl)-3-(2-methyl-1,2,3,4-tetrahydroisoquinolin-7-yl)acrylamide;  
 E-N-(4-Fluorophenyl)-3-(2-methyl-1,2,3,4-tetrahydroisoquinolin-7-yl)acrylamide;  
 E-N-(4-Methoxyphenyl)-3-(2-methyl-1,2,3,4-tetrahydroisoquinolin-7-yl)acrylamide;  
 15 E-N-(4-Carbomethoxyphenyl)-3-(2-methyl-1,2,3,4-tetrahydroisoquinolin-7-yl)acrylamide;  
 E-N-(4-Cyanophenyl)-3-(2-methyl-1,2,3,4-tetrahydroisoquinolin-7-yl)acrylamide;  
 E-N-(4-Nitrophenyl)-3-(2-methyl-1,2,3,4-tetrahydroisoquinolin-7-yl)acrylamide;  
 20 E-N-(4-Methylphenyl)-3-(2-methyl-1,2,3,4-tetrahydroisoquinolin-7-yl)acrylamide;  
 E-N-(3-Methoxy-5-trifluoromethylphenyl)-3-(2-methyl-1,2,3,4-tetrahydroisoquinolin-7-yl)acrylamide;  
 E-1-(3,4-Dihydro-1H-isoquinolin-2-yl)-3-(2-methyl-1,2,3,4-tetrahydroisoquinolin-7-yl)propenone;  
 25 E-1-(3,4-Dihydro-2H-quinolin-1-yl)-3-(2-methyl-1,2,3,4-tetrahydroisoquinolin-7-yl)propenone;  
 E-1-(3,3-Dimethyl-2,3-dihydroindol-1-yl)-3-(2-methyl-1,2,3,4-tetrahydroisoquinolin-7-yl)propenone, and;  
 E-1-(2,3-Dihydroindol-1-yl)-3-(2-methyl-1,2,3,4-tetrahydroisoquinolin-7-yl)propenone.  
 30

When synthesised, these compounds may be isolated in salt form, such as the hydrochloride or trifluoroacetate, and such salts also form part of this invention. Such salts may be used in preparing pharmaceutically acceptable salts. The compounds and their salts may be obtained as solvates, such as hydrates, and these also form part of this invention.

The above compounds and pharmaceutically acceptable salts thereof, especially the hydrochloride, and pharmaceutically acceptable solvates, especially hydrates, form a preferred aspect of the present invention.

The administration of such compounds to a mammal may be by way of oral, parenteral, sub-lingual, nasal, rectal or transdermal administration.

An amount effective to treat the disorders hereinbefore described depends on the usual factors such as the nature and severity of the disorders being treated and the weight of the mammal. However, a unit dose will normally contain 1 to 1000 mg, suitably 1 to 500 mg, for example an amount in the range of from 2 to 400 mg such as 2, 5, 10, 20, 30, 40, 50, 100, 200, 300 and 400 mg of the active compound. Unit doses will normally be administered once or more than once per day, for example 1, 2, 3, 4, 5 or 6 times a day, more usually 1 to 4 times a day, such that the total daily dose is normally in the range, for a 70 kg adult of 1 to 1000 mg, for example 1 to 500 mg, that is in the range of approximately 0.01 to 15 mg/kg/day, more usually 0.1 to 6 mg/kg/day, for example 1 to 6 mg/kg/day.

It is greatly preferred that the compound of formula (I) is administered in the form of a unit-dose composition, such as a unit dose oral, including sub-lingual, rectal, topical or parenteral (especially intravenous) composition.

Such compositions are prepared by admixture and are suitably adapted for oral or parenteral administration, and as such may be in the form of tablets, capsules, oral liquid preparations, powders, granules, lozenges, reconstitutable powders, injectable and infusable solutions or suspensions or suppositories. Orally administrable compositions are preferred, in particular shaped oral compositions, since they are more convenient for general use.

Tablets and capsules for oral administration are usually presented in a unit dose, and contain conventional excipients such as binding agents, fillers, diluents, tableting agents, lubricants, disintegrants, colorants, flavourings, and wetting agents. The tablets may be coated according to well known methods in the art. Suitable fillers for use include cellulose, mannitol, lactose and other similar agents. Suitable disintegrants include starch, polyvinylpyrrolidone and starch derivatives such as sodium starch glycollate. Suitable lubricants include, for example, magnesium stearate. Suitable pharmaceutically acceptable wetting agents include sodium lauryl sulphate.

These solid oral compositions may be prepared by conventional methods of blending, filling, tableting or the like. Repeated blending operations may be used to distribute the active agent throughout those compositions employing large quantities of fillers. Such operations are, of course, conventional in the art.

Oral liquid preparations may be in the form of, for example, aqueous or oily suspensions, solutions, emulsions, syrups, or elixirs, or may be presented as a dry product for reconstitution with water or other suitable vehicle before use. Such liquid preparations may contain conventional additives such as suspending agents, for example sorbitol, syrup, methyl cellulose, gelatin,

hydroxyethylcellulose, carboxymethyl cellulose, aluminium stearate gel or hydrogenated edible fats, emulsifying agents, for example lecithin, sorbitan monooleate, or acacia; non-aqueous vehicles (which may include edible oils), for example, almond oil, fractionated coconut oil, oily esters such as esters of  
5 glycerine, propylene glycol, or ethyl alcohol; preservatives, for example methyl or propyl *p*-hydroxybenzoate or sorbic acid, and if desired conventional flavouring or colouring agents. Oral formulations also include conventional sustained release formulations, such as tablets or granules having an enteric coating.

For parenteral administration, fluid unit dose forms are prepared  
10 containing the compound and a sterile vehicle. The compound, depending on the vehicle and the concentration, can be either suspended or dissolved. Parenteral solutions are normally prepared by dissolving the compound in a vehicle and filter sterilising before filling into a suitable vial or ampoule and sealing.  
Advantageously, adjuvants such as a local anaesthetic, preservatives and buffering  
15 agents are also dissolved in the vehicle. To enhance the stability, the composition can be frozen after filling into the vial and the water removed under vacuum.

Parenteral suspensions are prepared in substantially the same manner except that the compound is suspended in the vehicle instead of being dissolved and sterilised by exposure to ethylene oxide before suspending in the sterile  
20 vehicle. Advantageously, a surfactant or wetting agent is included in the composition to facilitate uniform distribution of the compound of the invention.

As is common practice, the compositions will usually be accompanied by written or printed directions for use in the medical treatment concerned.

Accordingly, the present invention further provides a pharmaceutical  
25 composition for use in the treatment and/or prevention of anxiety, mania, depression, panic disorders and/or aggression, disorders associated with a subarachnoid haemorrhage or neural shock, the effects associated with withdrawal from substances of abuse such as cocaine, nicotine, alcohol and benzodiazepines, disorders treatable and/or preventable with anti-convulsive agents, such as  
30 epilepsy including post-traumatic epilepsy, Parkinson's disease, psychosis, migraine, cerebral ischaemia, Alzheimer's disease and other degenerative diseases such as Huntingdon's chorea, schizophrenia, obsessive compulsive disorders (OCD), neurological deficits associated with AIDS, sleep disorders (including circadian rhythm disorders, insomnia & narcolepsy), tics (e.g. Gilles de la  
35 Tourette's syndrome), traumatic brain injury, tinnitus, neuralgia, especially trigeminal neuralgia, neuropathic pain, dental pain, cancer pain, inappropriate neuronal activity resulting in neurodysthesias in diseases such as diabetes, multiple sclerosis (MS) and motor neurone disease, ataxias, muscular rigidity (spasticity), temporomandibular joint dysfunction, and amyotrophic lateral


sclerosis (ALS) which comprises a compound of formula (I), or a pharmaceutically acceptable salt or solvate thereof, and a pharmaceutically acceptable carrier.

- 5 The present invention also provides a method of treatment and/or prevention of anxiety, mania, depression, panic disorders and/or aggression, disorders associated with a subarachnoid haemorrhage or neural shock, the effects associated with withdrawal from substances of abuse such as cocaine, nicotine, alcohol and benzodiazepines, disorders treatable and/or preventable with anti-convulsive agents, such as epilepsy including post-traumatic epilepsy, Parkinson's disease, psychosis, migraine, cerebral ischaemia, Alzheimer's disease and other  
10 degenerative diseases such as Huntingdon's chorea, schizophrenia, obsessive compulsive disorders (OCD), neurological deficits associated with AIDS, sleep disorders (including circadian rhythm disorders, insomnia & narcolepsy), tics (e.g. Gilles de la Tourette's syndrome), traumatic brain injury, tinnitus, neuralgia, especially trigeminal neuralgia, neuropathic pain, dental pain, cancer pain,  
15 inappropriate neuronal activity resulting in neurodysthesias in diseases such as diabetes, multiple sclerosis (MS) and motor neurone disease, ataxias, muscular rigidity (spasticity), temporomandibular joint dysfunction, and amyotrophic lateral sclerosis (ALS) comprising administering to the sufferer in need thereof an  
20 effective or prophylactic amount of a compound of formula (I), or a pharmaceutically acceptable salt or solvate thereof.


- In a further aspect the invention provides the use of a compound of formula (I), or a pharmaceutically acceptable salt or solvate thereof, for the manufacture of a medicament for the treatment and/or prevention of anxiety,  
25 mania, depression, panic disorders and/or aggression, disorders associated with a subarachnoid haemorrhage or neural shock, the effects associated with withdrawal from substances of abuse such as cocaine, nicotine, alcohol and benzodiazepines, disorders treatable and/or preventable with anti-convulsive agents, such as epilepsy including post-traumatic epilepsy, Parkinson's disease, psychosis,  
30 migraine, cerebral ischaemia, Alzheimer's disease and other degenerative diseases such as Huntingdon's chorea, schizophrenia, obsessive compulsive disorders (OCD), neurological deficits associated with AIDS, sleep disorders (including circadian rhythm disorders, insomnia & narcolepsy), tics (e.g. Gilles de la Tourette's syndrome), traumatic brain injury, tinnitus, neuralgia, especially  
35 trigeminal neuralgia, neuropathic pain, dental pain, cancer pain, inappropriate neuronal activity resulting in neurodysthesias in diseases such as diabetes, multiple sclerosis (MS) and motor neurone disease, ataxias, muscular rigidity (spasticity), temporomandibular joint dysfunction, and amyotrophic lateral sclerosis (ALS).

In a further aspect the invention provides the use of a compound of formula (I), or a pharmaceutically acceptable salt or solvate, thereof as a therapeutic agent, in particular for the treatment and/or prevention of anxiety, mania, depression, panic disorders and/or aggression, disorders associated with a subarachnoid haemorrhage or neural shock, the effects associated with withdrawal from substances of abuse such as cocaine, nicotine, alcohol and benzodiazepines, disorders treatable and/or preventable with anti-convulsive agents, such as epilepsy including post-traumatic epilepsy, Parkinson's disease, psychosis, migraine, cerebral ischaemia, Alzheimer's disease and other degenerative diseases such as Huntingdon's chorea, schizophrenia, obsessive compulsive disorders (OCD), neurological deficits associated with AIDS, sleep disorders (including circadian rhythm disorders, insomnia & narcolepsy), tics (e.g. Gilles de la Tourette's syndrome), traumatic brain injury, tinnitus, neuralgia, especially trigeminal neuralgia, neuropathic pain, dental pain, cancer pain, inappropriate neuronal activity resulting in neurodysthesis in diseases such as diabetes, multiple sclerosis (MS) and motor neurone disease, ataxias, muscular rigidity (spasticity), temporomandibular joint dysfunction, and amyotrophic lateral sclerosis (ALS).


The present invention also provides a process for the preparation of compounds of formula (I), which comprises  
 (a). for compounds of formula (I) in which P is -NH- and Q is -CR<sup>1</sup>=CH-, reacting a compound of formula (II)




with a compound of formula (III)



or,  
 (b) for compounds of formula (I) in which P is -CH=CH- and Q is -NR<sup>1</sup>-, reacting a compound of formula (IV)



with a compound of formula (V)



where  $R^{1A}$ ,  $R^{2A}$ ,  $R^{3A}$ ,  $R^{7A}$ ,  $R^{8A}$ ,  $R^{9A}$ , and  $R^{10A}$  are independently  $R^1$ ,  $R^2$ ,  $R^3$ ,  $R^7$ ,  $R^8$ ,  $R^9$ , and  $R^{10}$  as defined for formula (I) or a group or groups convertible thereto; Z, X and Y are as defined for formula (I); and L is OH or a halogen;

and where required converting an  $R^{1A}$ ,  $R^{2A}$ ,  $R^{3A}$ ,  $R^{7A}$ ,  $R^{8A}$ ,  $R^{9A}$ , or  $R^{10A}$  group to an  $R^1$ ,  $R^2$ ,  $R^3$ ,  $R^7$ ,  $R^8$ ,  $R^9$ , or  $R^{10}$  group;

converting one  $R^1$ ,  $R^2$ ,  $R^3$ ,  $R^7$ ,  $R^8$ ,  $R^9$ , or  $R^{10}$  group to another  $R^1$ ,  $R^2$ ,  $R^3$ ,  $R^7$ ,  $R^8$ ,  $R^9$ , or  $R^{10}$  group;


converting a salt product to the free base or another pharmaceutically acceptable salt, or converting a free base product to a pharmaceutically acceptable salt.

Conventional conditions for condensation of amines with carboxylic acids or active derivatives thereof, such as acid chlorides, may be used. For example the amides and acids may be reacted in the presence of a mixture of ethyl(dimethylaminopropyl)-carbodiimide/hydroxybenzotriazole in a suitable solvent such as dimethyl formamide, and amines and acid chlorides may be reacted together in a suitable solvent such as ethyl acetate or tetrahydrofuran. Alternatively the acid may be treated in solution with oxalyl chloride and then reacted with the amine or its hydrochloride.


Reaction of a compound of formula (III) or (V) which is an acid chloride ( $L=Cl$ ) in the absence of a base such as triethylamine will lead to formation of the hydrochloride salt of the compound of formula (I). In the presence of a base such as triethylamine the free base will be prepared. Hydrochloride salts can also be obtained by passing HCl gas into a solution of the free base, or adding a solution of HCl in ether.

Conversions of an  $R^{1A}$ ,  $R^{2A}$ ,  $R^{3A}$ ,  $R^{7A}$ ,  $R^{8A}$ ,  $R^{9A}$ , or  $R^{10A}$  group to an  $R^1$ ,  $R^2$ ,  $R^3$ ,  $R^7$ ,  $R^8$ ,  $R^9$ , or  $R^{10}$  group typically arise when a protecting group is needed during the above coupling reaction or during the preparation of the reactants by the procedures described below. Interconversion of one  $R^1$ ,  $R^2$ ,  $R^3$ ,  $R^7$ ,  $R^8$ ,  $R^9$ , or  $R^{10}$  group to another typically arises when one compound of formula (I) is used as the precursor of another compound of formula (I) or when it is easier to introduce a more complex or reactive substituent at the end of a synthetic sequence.


Compounds of formula (II) in which X is N (i.e. tetrahydronaphthyridines) may be prepared starting from a dinitro-1-methyl-2-pyridone compound of formula (VI)



by reaction with a 4-piperidone compound of formula (VII)



in a solution of ammonia in a suitable solvent such as methanol, to obtain a compound of formula (VIII) using a procedure similar to that of S Takada *et al*, J Med Chem, 1996, 39, 2844.







Compounds of formula (VIII) may be converted to compounds of formula (II) wherein X is nitrogen and Y is hydrogen by hydrogenation or reduction of the nitro group. For example, a compound of formula (VIII) may be hydrogenated by treatment with hydrogen in a suitable solvent such as methanol in the presence of a palladium/carbon catalyst. Alternatively, a compound of formula (VIII) may be reduced with stannous chloride in concentrated hydrochloric acid in a suitable solvent such as ethanol.

Compounds of formula (VI) may be prepared using the procedure of E. Matsumura, M. Ariga and Y. Tohda, Bull. Chem. Soc. Japan, 52 (8), 2413-2419 (1979).

Compounds of formula (II) in which X is CH and R<sup>3A</sup>, R<sup>7A</sup>, R<sup>8A</sup>, R<sup>9A</sup>, and R<sup>10A</sup> are hydrogen (i.e. tetrahydroisoquinolines) may be prepared from the corresponding unsaturated compound of formula (IX)




by reaction with a compound R<sup>3A</sup>M where M is a leaving group such as halogen, especially iodo, or tosylate to obtain an intermediate of formula (X)




which can be reduced, for example using sodium borohydride, to the compound of formula (II) wherein R<sup>7A</sup>, R<sup>8A</sup>, R<sup>9A</sup>, and R<sup>10A</sup> are hydrogen. Alternatively the compound of formula (X) can be hydrogenated, for example using hydrogen at 50psi in a solution of acetic/sulphuric acid with a platinum oxide catalyst.

Another route is from a precursor of formula (XI)



which can be reacted with R<sup>3A</sup>M, preferably as a tosylate, to obtain the intermediate of formula (XII)



(XII)


which can then be hydrogenated under the conditions previously described to  
 5 prepare the compound of formula (II) wherein X is CH and R<sup>7A</sup>, R<sup>8A</sup>, R<sup>9A</sup>, and R<sup>10A</sup> are hydrogen.

Compounds of formulae (IX) and (XI) and the reagents used are commercially available, or can be prepared from commercially available materials using conventional procedures described in the literature.


10 Alternatively, a compound of formula (II) wherein R<sup>7A</sup>, R<sup>8A</sup>, R<sup>9A</sup>, and R<sup>10A</sup> are hydrogen may be prepared directly from the corresponding nitro compound by catalytic hydrogenation. More specifically 7-amino-tetrahydroisoquinolines may be prepared by the procedure of G E Stokker, Tet. Lett. 1996, 37, 5453.

15 When R<sup>3A</sup> is hydrogen, the compound of formula (II) wherein R<sup>7A</sup>, R<sup>8A</sup>, R<sup>9A</sup>, or R<sup>10A</sup> are hydrogen can be obtained by direct hydrogenation of the compounds of formula (IX) or (XI), using the reagents already described. The NH group may be protected conventionally, for example by making R<sup>3A</sup> *t*-butoxycarbonyl prior to coupling and then deprotecting R<sup>3A</sup> under standard  
 20 conditions, for example using trifluoroacetic acid/methylene chloride.

Compounds of Formula (IV) may be prepared by initially reacting a compound of formula (XII) or formula (XIII)



(XII)



(XIII)

25 where Hal is a halogen, especially bromine, with an acrylate ester such as ethyl acrylate in conventional conditions. For example the reactants may be heated in the presence of palladium acetate and triethylamine in a suitable solvent such as acetonitrile. This reaction produces the corresponding ester of the L = OH  
 30

compound of formula (IV). Deesterification, for example by treatment with sodium or potassium hydroxide, gives the acid ( $L = OH$ ) of formula (IV). The acid of formula (IV) can be reacted with the amine of formula (V) under conditions mentioned above for reaction of formulae (II) and (III), or converted to the acid chloride, for example by treatment with carbonyl chloride, and then reacted with the amine.

In the reaction of formulae (IV) and (V), the group  $R^{3A}$  may be a desired substituent or a protecting group such as carboxylic acid *tert*-butyl ester. The protecting group may be removed at the end of the coupling to provide a compound in which  $R^3$  is H, or to provide a site for introduction of other  $R^3$  groups.

The halo compound (XIII) can be prepared by conventional means from commercial starting materials.

Compounds of formula (XII) are novel and form a further aspect of the present invention.

The aromatic amine compounds of formula (V), typically substituted phenylamines or bicyclic heterocycles such as tetrahydro(iso)quinolines and dihydroindolines, and the cinnamic acid derivatives of formula (III) are also commercially available or obtainable by conventional manipulation of substituents on aromatic acids and amines that are commercially available.

The above described procedures have been based on compounds in which Y is H. Compounds in which Y is a halogen may be prepared by reacting a compound of formula (II), or one of the above described precursors thereof (having an  $R^{3A}$  acting protecting group or an  $R^3$  substituent other than hydrogen) with a N-halo-succinimide in a suitable solvent such as acetonitrile, or N-chloromorpholine in a suitable solvent such as acetic acid for compounds where Y is chloro.

When  $R^{3A}$  is a protecting group then desired  $R^3$  substituents can be introduced into compounds of formula (II) or (IV) by removal of the protecting group followed by conventional N-substitutions, such as reaction with an appropriate aldehyde in the presence of a suitable reducing agent such as sodium borohydride.

Interconversions where Y is halogen, especially bromo or iodo, into intermediates of formula (II) where Y is alkyl can be carried out using a tetraalkyltin reagent in the presence of a suitable catalyst such as *bis* (triphenylphosphine) Pd (II) dichloride in a suitable solvent such as dimethylformamide at elevated temperature, optionally under argon. Alternatively, compounds of formula (I) where the  $R^2$  substituent is other than halogen and Y is bromo or iodo can also be converted into compounds of formula

(I) where Y is alkyl using an appropriate tetraalkyltin reagent. Other procedures for the interconversion of Y = halogen into Y = alkyl can be found in J. K. Stille, J. Org. Chem., 1990, 55, 3019.

- 5 Methods for the preparations of intermediates to other compounds of formula (II) where R<sup>7</sup>, R<sup>8</sup>, R<sup>9</sup>, R<sup>10</sup>, R<sup>11</sup>, R<sup>12</sup>, are alkyl can be found in WO98/41507; T.G.N.Watson., J. Org. Chem., 1998, 63, 406 [for R<sup>11</sup>, R<sup>12</sup>, as alkyl] and H. Takechi *et al.*, Synthesis 1992, 778 [for R<sup>7</sup>, R<sup>8</sup>, as alkyl].

- 10 *The preparation of compounds of formulae (II) and (IV) is illustrated by the following Descriptions; the preparation of compounds of this invention is illustrated by the following Examples. The utility of compounds of this invention is shown by the Pharmacological Data that follow the Examples. In the Descriptions and Examples, previously made compounds are referred to as, for example, "D1c" - meaning a compound made by Description 1c - and "E19rc" - a compound made as in Example 19rc)*

15

#### Description 1c

##### N-2-(4-Nitrophenyl)ethyl-trifluoroacetamide

- A solution of trifluoroacetic anhydride (10.6ml) in dichloromethane (100ml) was added dropwise to a stirred solution of 2,6- lutidine (17.44ml) and 4-  
20 nitrophenethylamine hydrochloride (15.2g; 75 mmol) at 0°C. The mixture was stirred at 25°C overnight under argon and then washed with dilute citric acid (x2), brine and dried over Na<sub>2</sub>SO<sub>4</sub>. The material in the organic phase gave the title compound as a pale yellow solid (19.04g).

- 25 **Description 2c**

##### 7-Nitro-1,2,3,4-tetrahydro-2-trifluoroacetyl-isoquinoline

- The nitro compound **D1c** (2.26g; 9.15 mmol) and paraformaldehyde (0.45g; 14.4 mmol) in acetic acid (10ml) and conc. H<sub>2</sub>SO<sub>4</sub> (15ml) were stirred at 25°C for 20h according to the procedure of G.E. Stokker., Tet. Lett., 1996, 37, 5453. Work up  
30 afforded the title compound as a white solid (2.17g).

<sup>1</sup>H NMR (CDCl<sub>3</sub>) δ: 3.10 (2H, m), 3.92 (2H, m), 4.85 + 4.92 (2H, 2xs), 7.38 (1H, t), 8.10 (2H, m); <sup>m/z</sup> (EI): 274 (M<sup>+</sup>)

#### Description 3c

- 35 **7-Nitro-1,2,3,4-tetrahydroisoquinoline**

The trifluoroacetamide **D2c** (17.22g; 63 mmol) was hydrolysed at room temperature using a solution of potassium carbonate (46.6g) in 10% aqueous methanol (660ml). Work-up with dichloromethane gave the title compound (11g).

**Description 4c****2-Methyl-7-nitro-1,2,3,4-tetrahydroisoquinoline**

- The amine D3c (2.08g; 11.7 mmol) was treated with 88% formic acid (3.45ml) and 37% aqueous formaldehyde (5.88ml) at 80°C for 2h according to the procedure of G.M. Carrera and D.S. Garvey, J. Het. Chem., 1992, 29, 847. Basification with 10% NaOH followed by work-up with ethyl acetate afforded an orange gum (2.3g). Chromatography on Kieselgel 60 in 0-3% methanol - ethyl acetate gave the title compound as an orange solid (1.7g).
- MS  $m/z$  (CI): 193 (MH<sup>+</sup>).

**Description 5c****7-Amino-2-methyl-1,2,3,4-tetrahydroisoquinoline**

- The 7-nitro compound D4c (0.25g; 1.3 mmol) in methanol (40ml) was hydrogenated over 10% palladium on carbon (100mg) at atmospheric pressure overnight. The catalyst was removed by filtration through a pad of Kieselguhr and evaporation *in vacuo* gave the title compound as a white solid (213mg).
- MS  $m/z$  (CI): 163 (MH<sup>+</sup>)

**Description 6c****7-Amino-2-(*t*-butoxycarbonyl)-1,2,3,4-tetrahydroisoquinoline**

- The title compound was prepared from the compound of D3c using di *t*-butyl dicarbonate in 10% aqueous hydroxide in dioxan at 25°C followed by catalytic hydrogenation according to the procedure of Description 5c.

**Description 7c****7-Amino-1,2,3,4-tetrahydro-2-trifluoroacetyl-isoquinoline**

- The 7-nitro compound D2c (0.99g; 3.6 mmol) in ethanol (50ml) was hydrogenated over 10% palladium on carbon (450mg) at atmospheric pressure for 4h. The catalyst was removed by filtration through a pad of Celite and evaporation *in vacuo* gave the title compound as a white solid (840mg).
- <sup>1</sup>H NMR (250MHz, CDCl<sub>3</sub>) δ: 2.84 (2H, t), 3.23 (2H, br s), 3.82 (2H, m), 4.66 (2H, d, restricted rotation around C-1), 6.47 (1H, m), 6.57 (1H, m), 6.96 (1H, m)

**Description 8c****6-Methyl-3-nitro-5,6,7,8-tetrahydro[1,6]naphthyridine**

- 3,5-Dinitro-1-methyl-2-pyridone (5.97g; 30 mmol) was treated with 1.22M ammonia in methanol (300ml) followed by 1-methyl-4-piperidone (3.73g, 33

mmol) and the mixture heated at 60° for 5h, then allowed to stand at ambient temp for 72h. Evaporated to dryness under reduced pressure and the orange/red residue triturated under a mixture of dichloromethane and diethyl ether, collected by filtration, washed with diethyl ether and dried in air. Chromatography through silica gel, eluting with ethyl acetate, gave the title compound as a red solid (3.4g, 59%);  $\nu_{\max}$  (CH<sub>2</sub>Cl<sub>2</sub>) 1530 and 1351 cm<sup>-1</sup>  
<sup>1</sup>H NMR (250MHz, CDCl<sub>3</sub>)  $\delta$ : 2.53 (3H, s), 2.85 (2H, t, J = 6 Hz), 3.18 (2H, t, J = 6 Hz), 3.69 (2H, s), 8.14 (1H, d, J = 2 Hz), 9.23 (1H, d, J = 2 Hz)

10 **Description 9c**

**3-Amino-6-methyl-5,6,7,8-tetrahydro[1,6]naphthyridine**

6-Methyl-3-nitro-5,6,7,8-tetrahydro[1,6]naphthyridine (2.72g, 1.41 mmol) was dissolved in methanol (100ml) and treated with 10% palladium on carbon (1.0g). The mixture was hydrogenated for 2h. The catalyst was removed by filtration through Celite, the filter bed washed with methanol and the filtrate evaporated to dryness under reduced pressure to give a yellow solid, which was triturated under diethyl ether and the solids collected by filtration, washed with diethyl ether and dried *in vacuo* (1.89g, 83%)

<sup>1</sup>H NMR (250MHz, CDCl<sub>3</sub>)  $\delta$ : 2.46 (3H, s), 2.75 (2H, t, J = 6 Hz), 2.95 (2H, t, J = 6 Hz), 3.50 (2H, s), 3.56 (2H, br s, exchangeable), 6.65 (1H, d, J = 2 Hz), 7.92 (1H, d, J = 2 Hz)

**Description 10c**

**5-Amino-2-methylisoquinolinium iodide**

25 To a solution of 5-aminoisoquinoline (14.4g, 100mmol) in acetone (300ml) was added iodomethane (14.4ml). The solution was briefly stirred and then allowed to stand for 2h. The yellow precipitate was then filtered, washed with acetone and dried to afford the title compound as a yellow solid (18.8g).

30 **Description 11c**

**5-Amino-2-methyl-1,2,3,4-tetrahydroisoquinoline**

Sodium borohydride (17.8g, 0.47mol) was added portionwise over 2h to an ice cold solution of 5-amino-2-methylisoquinolinium iodide (18.8g, 65mmol) in methanol (1.5L) and water (60ml). The mixture was then stirred at 25°C for 18h. and concentrated *in vacuo*. The residue was extracted into water and dichloromethane. The organic layer was dried (Na<sub>2</sub>SO<sub>4</sub>) and concentration *in vacuo* gave the title compound (8.87g).

**Description 12c**

**7-Amino-1,2,3,4-tetrahydro-2-trifluoroacetyl-isoquinoline**

The 7-nitro compound **D2c** (0.99g; 3.6 mmol) in ethanol (50ml) was hydrogenated over 10% palladium on carbon (450mg) at atmospheric pressure for 4h. The catalyst was removed by filtration through a pad of Celite and evaporation *in vacuo* gave the title compound as a white solid (840mg).

<sup>1</sup>H NMR (250MHz, CDCl<sub>3</sub>) δ: 2.84 (2H, t), 3.23 (2H, br s), 3.82 (2H, m), 4.66 (2H, d, restricted rotation around C-1), 6.47 (1H, m), 6.57 (1H, m), 6.96 (1H, m)

**Description 13c****7-Amino-8-chloro-1,2,3,4-tetrahydro-2-trifluoroacetyl-isoquinoline**

To a solution of amine **D12c** (1.00g) in acetonitrile (20ml) N-chlorosuccinimide (0.60g) was added and the solution stirred at room temperature for 6 days. The solution was diluted with ethyl acetate, washed with water and the organic phase dried (MgSO<sub>4</sub>) and solvent removed at reduced pressure. The residue was column chromatographed (silica gel, dichloromethane then 2% methanol/dichloromethane) to give 7-amino-8-chloro-1,2,3,4-tetrahydro-2-trifluoroacetyl-isoquinoline as a pale yellow solid (0.72g).

<sup>1</sup>H NMR (250MHz, CDCl<sub>3</sub>) δ: 2.85 (2H, m), 3.83 (2H, dt, restricted amide rotation), 4.76 (2H, s), 6.68 (1H, m) and 6.89 (1H, m).

**Description 14c****7-Amino-8-bromo-1,2,3,4-tetrahydro-2-trifluoroacetylisoquinoline**

The title compound (0.27g) was prepared from amine **D12c** (0.24g) and N-bromosuccinimide (0.20g) according to the method of Description 13c.

<sup>1</sup>H NMR (250MHz, CDCl<sub>3</sub>) δ: 2.85 (2H, m), 3.76 - 3.87 (2H, m, restricted amide rotation), 4.72 (2H, d due to restricted amide rotation), 6.68 (1H, m) and 6.93 (1H, m).

**Description 15c****5-Iodo-7-nitro-1,2,3,4-tetrahydroisoquinoline**

The nitro compound **D3c** (750mg; 3.9mmol) and N-iodosuccinimide (1.13g) in triflic acid (5ml) was stirred at 25°C overnight. The mixture was poured cautiously into saturated NaHCO<sub>3</sub> and then extracted into ether (2x). The combined organic extracts were washed with aqueous sodium thiosulfate, dried (MgSO<sub>4</sub>) and evaporation *in vacuo* gave a residue. Chromatography on Kieselgel 60 in 2% methanol - dichloromethane gave the title compound (650mg).

**Description 16c****5-Iodo-7-nitro-2-trifluoroacetyl-1,2,3,4-tetrahydroisoquinoline**

The title compound was prepared from **D15c** and trifluoroacetic anhydride using a procedure similar to that of Description 6.

MS m/z (API<sup>+</sup>): 401 (MH<sup>+</sup>; 45%).

5 **Description 17**

**5-Chloro-7-nitro-2-trifluoroacetyl-1,2,3,4-tetrahydroisoquinoline**

**D16c** (810mg) in dry DMF (15ml) was treated with copper (I) chloride (605mg) and heated at 125°C under argon for 18h. After cooling, the mixture was concentrated *in vacuo* and the residue partitioned between ethyl acetate and water.

10 The organic layer was washed with water (x 3), aqueous sodium thiosulfate, brine and dried (MgSO<sub>4</sub>). Evaporation *in vacuo* gave the title compound as a red gum (519mg).

<sup>1</sup>H NMR (CDCl<sub>3</sub>) δ: 3.09 (2H, m), 3.96 (2H, m), 4.85, 4.92 (2H, 2s, rotamers), 7.99 (1H, m), 8.20 (1H, m).

15

**Description 18c**

**7-Amino-5-chloro-2-trifluoroacetyl-1,2,3,4-tetrahydroisoquinoline**

A solution of the nitro compound **D17c** (2.14mmol) in ethanol (20ml) at 50°C was treated with a solution of tin (II) chloride (1.42g) in c. HCl (3ml). The resultant yellow solution was basified with 10% aqueous sodium hydroxide and the product extracted into dichloromethane. Flash chromatography on Kieselgel 60 (5% methanol - dichloromethane) gave the title compound.

20

<sup>1</sup>H NMR (CDCl<sub>3</sub>) δ: 2.84 (2H, m), 3.67 (2H, brs), 3.83 (2H, m), 4.61, 4.67 (2H, 2s, rotamers), 6.33 (1H, m), 6.65 (1H, m).

25

**Description 19c**

**7-Amino-5-bromo-2-trifluoroacetyl-1,2,3,4-tetrahydroisoquinoline**

The title compound was prepared from **D16c** and copper (II) bromide using a method similar to that of Description 10 followed by tin (II) chloride reduction according to the procedure used in Description 18.

30

<sup>1</sup>H NMR (CDCl<sub>3</sub>) δ: 2.86 (2H, m), 3.68 (2H, brs), 3.85 (2H, m), 4.62, 4.69 (2H, 2s, rotamers), 6.39 (1H, m), 6.85 (1H, m).

**Description 20c**

35 **7-Nitro-2,4,4-trimethyl-4H-isoquinoline-1,3-dione**

2,4,4-Trimethyl-4H-isoquinoline-1,3-dione (5g, 24.6mmol) [prepared according to H. Takechi *et al.*, Synthesis. 1992, 778] in concentrated sulfuric acid (50ml) at 0°C was treated with fuming nitric acid (2.5ml, dropwise) over 5 min and the



reaction warmed to 25°C. After stirring for 30 min at 25°C the reaction mixture was poured into ice water (100ml) and the organics extracted into dichloromethane (3x50ml). The combined extracts were dried (MgSO<sub>4</sub>) and evaporated *in vacuo* to give the title compound (5.31g, 86%).

- 5 <sup>1</sup>H NMR (250 MHz, CDCl<sub>3</sub>) δ: 1.70 (6H, s), 3.42 (3H, s), 7.69 (1H, d, J = 9 Hz), 8.46 (1H, dd, J = 9, 2 Hz), 9.07 (1H, d, J = 2 Hz); <sup>m/z</sup> (AP<sup>+</sup>): 249 (M+H)<sup>+</sup>

#### Description 21c

##### 7-Amino-2,4,4-trimethyl-4H-isoquinoline-1,3-dione

- 10 7-Nitro-2,4,4-trimethyl-4H-isoquinoline-1,3-dione (45g, 20mmol) was dissolved in a methanol (500ml)/dichloromethane (100ml) mixture and treated with 10% Pd/C (0.5g). The reaction mixture was hydrogenated for 2h before removal of the palladium catalyst by filtration through Celite. The filtrate was evaporated to dryness *in vacuo* to give the title compound (4.4g, quant).

- 15 <sup>1</sup>H NMR (250 MHz, CDCl<sub>3</sub>) δ: 1.58 (6H, s), 3.36 (3H, s), 3.83 (2H, brs), 6.95 (1H, dd, J = 6, 3 Hz), 7.24 (1H, d, J = 6 Hz), 7.48 (1H, d, J = 3 Hz); MS <sup>m/z</sup> (AP<sup>+</sup>): 219 (M+H)<sup>+</sup>

#### Description 22c

- 20 7-Amino-2,4,4-trimethyl-1,2,3,4-tetrahydroisoquinoline, hydrochloride

- 7-Amino-2,4,4-trimethyl-4H-isoquinoline-1,3-dione (4g, 18.3mmol) was dissolved in tetrahydrofuran (400ml) and heated at reflux (~61°C). Borane-tetrahydrofuran complex (88ml, 1M solution in THF) was added dropwise to the mixture and heating continued for a further 3 h. The cooled reaction (0°C) was treated with methanol (400ml) dropwise to destroy residual borane, followed by evaporation *in vacuo*. The resultant residue was heated at reflux in 3N HCl (400ml) for 30 min. The mixture was cooled to 0°C and treated with NaOH pellets until basic (pH 9). The free amine was extracted into dichloromethane (4x100ml) before drying over magnesium sulfate and evaporation *in vacuo*. The resulting light brown oil was dissolved in dichloromethane (50ml) and treated with hydrogen chloride (1M solution in ether) until acidic (pH 2). Solvent removal *in vacuo* followed by trituration with ether yielded the title compound as an off-white powder (3.3g, 79%).

- 30 <sup>1</sup>H NMR (free base 250 MHz, CDCl<sub>3</sub>) δ: 1.25 (6H, s), 2.37 (2H, s), 2.39 (3H, s), 3.43 (2H, s), 3.51 (2H, brs), 6.32 (1H, d, J = 2 Hz), 6.54 (1H, dd, J = 8, 2 Hz), 7.09 (1H, d, J = 8 Hz); <sup>m/z</sup> (AP<sup>+</sup>): 191 (M+H)<sup>+</sup>

#### Description 23c

##### 3,4-Dihydro-3,3-dimethyl-7-nitroisoquinoline

- To a stirred solution of potassium nitrate (2.53g) in sulfuric acid (14ml) at 0°C was added dropwise a solution of 3,4-dihydro-3,3-dimethyl isoquinoline (3.68g; 23mmol) [prepared according to the procedure of T.J.N.Watson, *J. Org. Chem.*, 1998, 63, 406] in sulfuric acid (13.5ml). The resultant solution was stirred at
- 5 room temperature for 1.5h and then heated to 60°C for 4.5h. The solution was then cooled to room temperature, and poured on to ice.; 0.880 ammonia was added until the solution was neutral, and the product was extracted into dichloromethane (x3). The combined organic phases were, dried over magnesium sulphate, and then evaporated *in vacuo* to afford the title compound (4.22g).
- 10 <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ: 1.27 (6H, s), 2.85 (2H, s), 7.34 (1H, d, J = 8 Hz), 8.17 (1H, d, J = 2 Hz), 8.23 (1H, dd, J = 8, 2 Hz), 8.33 (1H, s).

#### Description 24c

##### 3,3-Dimethyl-7-nitro-1,2,3,4-tetrahydroisoquinoline

- 15 Sodium borohydride (1.57g; 41.38mmol) was added portionwise to a solution of 3,4-dihydro-3,3-dimethyl-7-nitroisoquinoline (4.22g; 20.69mmol) in methanol (150ml). The resultant solution was stirred at room temperature for 2 h. The methanol was evaporated *in vacuo* and the residue partitioned between water and dichloromethane. The organic layer was dried (Na<sub>2</sub>SO<sub>4</sub>) and then evaporated *in*
- 20 *vacuo* to afford the title compound (3.81g).
- <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ: 1.20 (6H, s), 1.40 - 1.53 (1H, brs), 2.72 (2H, s), 4.14 (2H, s), 7.20 (1H, d, J = 8 Hz), 7.37 (1H, s), 7.98 (1H, dd, J = 8, 2 Hz).

#### Description 25c

##### 3,3-Dimethyl-7-nitro-1,2,3,4-tetrahydro-2-trifluoroacetylisoquinoline

- 25 A solution of 2,6-lutidine (2.29ml; 19.69mmol) and 3,3-dimethyl-7-nitro-1,2,3,4-tetrahydroisoquinoline (3.7g; 17.9mmol) in dichloromethane (150ml) was treated dropwise, with ice cooling, trifluoroacetic anhydride (2.53ml, 17.9mmol) in dichloromethane (50ml). The reaction was then allowed to warm to 25°C and
- 30 stirred for 18h. The resultant mixture was washed with 5M HCl, brine, dried (Na<sub>2</sub>SO<sub>4</sub>) and then evaporated *in vacuo* to afford the title compound (5.82g).
- <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ: 1.51 (6H, s), 2.97 (2H, s), 4.61 (2H, s), 7.43 (1H, d, J = 8 Hz), 8.12 (1H, d, J = 2 Hz), 8.24 (1H, dd, J = 8, 2 Hz).

#### 35 Description 26c

##### 7-Nitro-2,3,3-trimethyl-3,4-dihydroisoquinolinium iodide

- D24c (1.0g, 4.9mmol) was dissolved in acetone (100ml) and treated with iodomethane (1ml, 16mmol). The reaction was stirred at room temperature for

18h. The resultant precipitate was collected by filtration and dried; pale yellow powder (1.5g, 88%).

MS m/z (API<sup>+</sup>): 219 (M)<sup>+</sup>

5 **Description 27c**

**7-Nitro-2,3,3-trimethyl-1,2,3,4-tetrahydroisoquinoline**

D26c (200mg, 5.8mmol) was reduced with sodium borohydride (300mg; 7.9mmol) in a manner similar to that of Description 24c. Purification by chromatography eluting with a dichloromethane solution of ammonia in methanol (0.5% conc. NH<sub>3</sub>: 4.5% MeOH: 95% CH<sub>2</sub>Cl<sub>2</sub>) gave the title compound as a pale yellow oil (93mg, 73%).

<sup>1</sup>H NMR (CDCl<sub>3</sub>) δ: 1.10 (6H, s), 2.40 (3H, s), 2.78 (2H, s), 3.80 (2H, s), 7.21 (1H, d, J = 8 Hz), 7.90 (2H, m).

15 **Description 28c**

**7-Amino-2,3,3-trimethyl-1,2,3,4-tetrahydroisoquinoline**

The title compound was prepared from D27c using a method similar to that of Description 2c. For ease of handling the compound was converted into a monohydrochloride.

20 <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ: 1.07 (6H, s), 2.35 (3H, s), 2.59 (2H, s), 3.46 (2H, brs), 3.64 (2H, s), 6.37 (1H, d, J = 2 Hz), 6.50 (1H, dd, J = 8, 2 Hz), 6.84 (1H, d, J = 8 Hz).

**Description 29c**

**7-Amino-8-chloro-2,3,3-trimethyl-1,2,3,4-tetrahydroisoquinoline**

25 Chlorination of D28c (900mg; 4.74 mmol) with N-chloromorpholine (600mg; 4.90 mmol) in glacial acetic acid (30ml) for 30 min at 25°C followed by basic work-up with dichloromethane gave the title compound (700mg).

<sup>1</sup>H NMR (CDCl<sub>3</sub>) δ: 1.06 (6H, s), 2.40 (3H, s), 2.60 (2H, s), 3.67 (2H, s), 3.92, (2H, brs), 6.62 (1H, d, J = 8 Hz), 6.79 (1H, d, J = 8 Hz); m/z (API<sup>+</sup>): 225.1 (MH<sup>+</sup>); 30 100% expected isotope pattern)

**Description 30c**

**7-Amino-8-bromo-2,4,4-trimethyl-1,2,3,4-tetrahydroisoquinoline**

35 To a solution of 7-amino-2,4,4-trimethyl-1,2,3,4-tetrahydroisoquinoline from D22c (7g) in acetonitrile (200ml), was added N-bromo succinimide (7.21g) portionwise over 10 min. The reaction mixture was cooled in an ice/methanol bath to prevent any large exotherm and then stirred under argon for 45 min. The reaction was allowed to warm to room temperature, diluted with water and extracted with ethyl acetate. The combined organic extracts were washed with

brine, dried (MgSO<sub>4</sub>) and evaporated to dryness *in vacuo* to afford a brown solid which was purified using dry flash column chromatography eluting with ethyl acetate. Combination of appropriate fractions afforded the title compound as an orange gum (3.95g).

- 5 <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ: 1.26 (6H, s), 2.33 (2H, s), 2.45 (3H, s), 3.49 (2H, s), 4.00 (2H, s), 6.67 (1H, d), 7.08 (1H, d).

#### Description 31c

##### 7-Amino-8-ethyl-2,4,4-trimethyl-1,2,3,4-tetrahydroisoquinoline

- 10 A solution of D30c (3.95g) and lithium chloride (1.87g) in dry dimethylformamide (120ml) was treated with tetraethyl tin (5.81ml) followed by a catalytic amount of *bis* (triphenylphosphine) palladium (II) dichloride (350mg). The reaction mixture was then stirred under argon at 120°C overnight. After cooling, the solvent was removed *in vacuo* and the residual oil was dissolved in dichloromethane and
- 15 filtered through Celite, washing with dichloromethane. The organic phase was evaporated *in vacuo* to afford a dark orange oil which was purified using dry flash column chromatography eluting with ethyl acetate. Combination of appropriate fractions gave the title compound as a yellow gum (1.6g).

- 20 <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ: 1.14 (3H, t), 1.27 (6H, d), 2.33 (2H, s), 2.47 (5H, m), 3.51 (2H, s), 6.60 (1H, d), 7.02 (1H, d).

#### Description 32c

##### 1,2-Dimethyl-3,4-dihydroisoquinolinium iodide

- 25 1-Methyl-3,4-dihydroisoquinoline (780mg) was dissolved in acetone (7ml) and iodomethane (0.38ml) added. The solution was allowed to stand overnight at room temperature. The product was obtained as pale yellow crystals (1.4g).
- <sup>1</sup>H NMR (250MHz, d<sub>6</sub>DMSO) δ: 2.66 (3H, s), 2.99 (2H, t, J = 7.5 Hz), 3.56 (3H, s), 3.88 (2H, t, J = 7.5 Hz), 7.36 (2H, m), 7.60 (1H, t, J = 7.5 Hz), 7.94 (1H, d, J = 7.5 Hz).

30

#### Description 33c

##### 1,1,2-Trimethyl-1,2,3,4-tetrahydroisoquinoline

- Methyl magnesium bromide (4.5ml, 3M in Et<sub>2</sub>O) was added to a stirred suspension of 1,2-dimethyl-3,4-dihydroisoquinolinium iodide (1.3g) in dry THF
- 35 (20ml) at -70°C under argon. After 1h, the mixture was allowed to warm slowly to room temperature and stirred overnight. The mixture was quenched by cautious addition of water and extracted with ethyl acetate. The extract was washed with water, brine, dried (MgSO<sub>4</sub>) and evaporated *in vacuo* to give the title compound as a pink oil (730mg).

<sup>1</sup>H NMR (250MHz, CDCl<sub>3</sub>) δ: 1.40 (6H, s), 2.44 (3H, s), 2.87 (4H, s), 7.15 (4H, m).

#### Description 34c

##### 5 1,1,2-Trimethyl-7-nitro-1,2,3,4-tetrahydroisoquinoline

The amine D33c (620mg) was converted into the sulfate salt and added to an ice-cooled solution of potassium nitrate (420mg) in conc. H<sub>2</sub>SO<sub>4</sub> (5ml). When the addition was complete the ice bath was removed and the mixture stirred overnight at room temperature. The mixture was poured onto ice, made basic with conc. aq. ammonia and extraction with dichloromethane yielded an oil which was purified by chromatography eluting with dichloromethane: methanol: ammonia. The product was obtained as a red oil (430mg) [predominantly the desired 7-nitro derivative].

10 <sup>1</sup>H NMR (250MHz, CDCl<sub>3</sub>) δ: 1.45 (6H, s), 2.45 (3H, s), 2.92 (4H, m), 7.20 (1H, d, J = 8 Hz), 7.95 (1H, dd, J = 8, 2 Hz), 8.15 (1H, d, J = 2 Hz).

#### Description 35c

##### 7-Amino-3,4-dihydroisoquinoline

7-Nitro-3,4-dihydroisoquinoline (0.60g, 3.4mmol) [prepared according to the procedure of A.P. Venkov *et al*, Syn. Commun., 1996 26 127] was dissolved in ethanol (100ml) and heated to 60°C. This hot solution was treated with a solution of tin (II) chloride dihydrate (3.08g, 13.7mmol) in conc. HCl (10ml). The resultant mixture was heated at 60° for 1h. Upon cooling, the reaction mixture was poured into water (100ml) and basified (pH 9) with KOH pellets, liberating an oily residue. This residue was extracted into dichloromethane and dried over magnesium sulfate. Purification by chromatography through silica gel, eluting with (0.5% conc. ammonia : 4.5% methanol : 95% dichloromethane) yielded the title compound as a dark yellow oil (0.44g, 88%).

25 <sup>1</sup>H NMR (250MHz, CDCl<sub>3</sub>) δ: 2.63 (2H, t, J = 7 Hz), 3.67 (2H, brs), 3.73 (2H, m, J = 7, 2 Hz), 6.62 (1H, d, J = 2 Hz), 6.70 (1H, dd, J = 8, 2 Hz), 6.95 (1H, d, J = 8 Hz), 8.24 (1H, s).

#### Description 36c

##### 7-Amino-2-methyl-3,4-dihydroisoquinolinium iodide

35 7-Amino-3,4-dihydroisoquinoline (0.40g, 2.74mmol) in acetone (125ml) was treated with iodomethane (0.50ml, 8.03mmol) and left stirring at room temperature for 18 h. The resultant yellow precipitate was collected by filtration and dried *in vacuo* at ambient temperature (0.73g, 92%).

MS <sup>m/z</sup> (API<sup>+</sup>): 161 (M)<sup>+</sup>

**Description 37c****(±) 7-Amino-1,2-dimethyl-tetrahydroisoquinoline**

(±) 7-Amino-2-methyl-3,4-dihydroisoquinolinium iodide (0.50g, 1.7mmol) was  
5 suspended in anhydrous tetrahydrofuran (50ml) and cooled to -78°C. The cooled  
solution was treated with methyl magnesium chloride (2.14ml of a 3M solution in  
THF, 6.96mmol), added as a single portion. The reaction was allowed to reach  
room temperature over 18 h before being poured into water (50ml). The organic  
solvent was removed *in vacuo* and the organic product extracted into  
10 dichloromethane. Drying over magnesium sulfate and evaporation *in vacuo*  
furnished the title compound as a pale yellow oil (0.3g, 98%). For ease of  
handling the product was converted into a monohydrochloride.  
<sup>1</sup>H NMR (250MHz, CDCl<sub>3</sub>) δ: 1.37 (3H, d, J = 7 Hz), 2.46 (3H, s), 2.54 - 2.83  
(3H, m), 3.00 (1H, m), 3.50 (3H, m), 6.45 (1H, d, J = 2 Hz), 6.51 (1H, dd, J = 8, 2  
15 Hz), 6.88 (1H, d, J = 8 Hz).

**Description 38c****4,4-Dimethyl-7-nitro-3,4-dihydroisoquinoline**

The title compound was prepared in a manner to that described in Description 35c.  
20 MS m/z (AP<sup>+</sup>): 205 (MH<sup>+</sup>; 100%).

**Description 39c****2,4,4-Trimethyl-7-nitro-3,4-dihydroisoquinolinium iodide.**

The title compound was prepared in a manner to that described in Description 36c.  
25 MS m/z (AP<sup>+</sup>): 219 (MH<sup>+</sup>; 100%).

**Description 40c****7-Nitro-1,2,4,4-tetramethyl-1,2,3,4-tetrahydroisoquinoline**

D39c (1.5g, 4.3mmol) in THF (50ml) was stirred under argon and dimethyl zinc in  
30 toluene (3.3ml, 2M solution) added with rapid stirring at 0°C. The mixture was  
allowed to warm to room temperature over 1h, quenched with satd. ammonium  
chloride and concentrated *in vacuo*. Work-up with dichloromethane gave the title  
compound (0.9g, 90%).  
MS m/z (AP<sup>+</sup>): 235 (MH<sup>+</sup>; 100%).  
35

**Description 41c****7-Amino-1,2,4,4-tetramethyl-1,2,3,4-tetrahydroisoquinoline**

Prepared from D40c in a manner similar to that of Description 2c.

<sup>1</sup>H NMR (CDCl<sub>3</sub>) δ: 1.34 (3H, s), 1.35 (3H, s), 1.49 (3H, d, J = 7 Hz), 2.55 - 2.66 (1H, m), 2.62 (3H, brs), 2.89 (1H, m), 3.50 (2H, brs), 3.77 - 3.90 (1H, m), 6.47 (1H, d, J = 2 Hz), 6.59 (1H, dd, J = 8, 2 Hz), 7.11 (1H, d, J = 8 Hz).

5    **Description 42c**

**7-Amino-8-methyl-2,4,4-trimethyl-1,2,3,4-tetrahydroisoquinoline**

A solution of D30c and lithium chloride in dry dimethylformamide was treated with tetramethyl tin in a manner similar to that of Description 31c.

10   **Description 43c**

**7-Amino-8-chloro-2,4,4-trimethyl-1,2,3,4-tetrahydroisoquinoline**

The title compound was prepared from D22c and N-chloromorpholine using a procedure similar to that of Description 29c.

15   **Description 44c**

**7-Amino-8-chloro-4,4-dimethyl-2-trifluoroacetyl-1,2,3,4-tetrahydroisoquinoline**

<sup>1</sup>H NMR (250MHz, CDCl<sub>3</sub>) δ: 1.27 (6H, s), 3.50, 3.63 (2H, 2s, rotamers), 4.05 (2H, brs), 4.76 (2H, s), 6.73 (1H, m), 7.07 (1H, m).

20

**Description 45c**

**1,3,3-Trimethylpiperidin-4-one**

The title compound was prepared according to the procedure of Katvalyan *et al.*, Bull. Acad. Sci. USSR (Engl) 1968, 2436.

25   b.p 70 °C at 16mm Hg; <sup>m/z</sup> (API<sup>+</sup>): 142.1 (MH<sup>+</sup>)

**Description 46c**

**3-Nitro-5,6,7,8-tetrahydro-6,8,8-trimethyl[1,6]naphthyridine**

30   3,5-Dinitro-1-methylpyridin-2-one [prepared by the method of E. Matsumura, M. Ariga and Y. Tohda, Bull. Chem. Soc. Japan, 1979, 52, 2413-2419] (2g; 10mmol) was suspended in MeOH (50ml) and treated with 0.88 aq. ammonia (10ml; 157mmol). 1,3,3-Trimethylpiperidin-4-one (1.7g; 12mmol) was added and the mixture heated at 70°C for 5h. The mixture was cooled to room temperature then evaporated to dryness *in vacuo*. The residue was digested with dichloromethane  
35   (2x50ml) and the hot solution decanted from the red gum. The extracts were combined, evaporated to dryness *in vacuo* and the residue purified by chromatography on SiO<sub>2</sub>, with 50% ethyl acetate : 60-80 °C petroleum to give the title compound as a yellow oil, which solidified on standing (1.05g; 48%).

<sup>1</sup>H NMR (250 MHz; CDCl<sub>3</sub>) δ: 1.38 (6H, s), 2.47 (3H, s), 2.55 (2H, s), 3.64 (2H, s), 8.09 (1H, d, J = 3 Hz), 9.25 (1H, d, J = 3 Hz); <sup>m/z</sup> (API<sup>+</sup>): 222.1 (MH<sup>+</sup>)

#### Description 47c

##### 5 3-Amino-5,6,7,8-tetrahydro-6,8,8-trimethyl[1,6]naphthyridine

The product from D46c (930mg; 4.20mmol) was dissolved in MeOH (30ml) and the mixture hydrogenated in a manner similar to that of Description 2 to give the title compound (795mg; 84%).

<sup>1</sup>H NMR (250 MHz; CD<sub>3</sub>OD) δ: 1.73 - 1.99 (2H, m), 2.34 - 2.55 (5H, m), 2.63 (1H, d, J = 17 Hz), 3.29 and 3.36 (1H, dd, J = 17, 5 Hz), 3.66 - 3.71 (1H, m), 3.99 (1H, d, J = 6 Hz), 6.95 (1H, d, J = 3 Hz), 7.95 (1H, d, J = 3 Hz).

#### Example 1c

##### E-N-(2-Methyl-1,2,3,4-tetrahydroisoquinolin-7-yl)-3-nitrocinnamide

15 3-Nitrocinnamic acid (195mg; 1.0 mmol), ethyldimethylaminopropyl carbodiimide (194mg; 1.3 mmol) and 1-hydroxybenzotriazole (136mg; 1.0 mmol) in dry DMF (12ml) was stirred at room temperature for 30 min. A solution of the N-methyl amine D5c (164mg; 1.0 mmol) in dichloromethane (5ml) was added and the mixture kept at room temperature overnight. The resultant cream precipitate  
20 was removed by filtration and the residue washed well with ether:hexane. The residue was dried *in vacuo* and gave the title compound as an off white solid (0.5g).

<sup>1</sup>H NMR (250MHz, d<sup>6</sup> DMSO) δ: 2.35 (2H, br, overlapping), 2.74 (3H, s), 2.90 (2H, br), 4.20 (2H, br), 6.90 (1H, d, J = 16 Hz), 7.07 (1H, d, J = 8Hz), 7.39 (1H, dd), 7.50 (1H, d, J = 16Hz), 7.60 (1H, t), 7.94 (1H, d, J = 8Hz), 8.10 (1H, m), 8.32 (1H, narrow t);

MS <sup>m/z</sup> (API): 338 (MH<sup>+</sup>; 100%)

#### Example 2c

##### 30 E-N-(2-Methyl-1,2,3,4-tetrahydroisoquinolin-7-yl)-3-trifluoromethylcinnamide

3-Trifluoromethylcinnamoyl chloride (234mg; 1.0mmol) was added to a stirred solution of amine D5c (162mg; 1.0mmol) in dichloromethane (25ml) containing dry triethylamine (0.3ml). The mixture was kept at room temperature overnight  
35 and work-up similar to that for Example 1c, followed by flash chromatography on Kieselgel 60 (10% methanol:ethyl acetate) gave the title compound as a buff powder (200mg; 55%).



<sup>1</sup>H NMR (250MHz, CDCl<sub>3</sub>) δ: 2.46 (3H, s), 2.69 (2H, t), 2.89 (2H, t), 3.56 (2H, s), 6.62 (1H, d, J = 16 Hz), 7.08 (1H, d, J = 6.6Hz), 7.30 (1H, m), 7.40 (1H, brs), 7.52 (1H, t), 7.64 (3H, m), 7.75 (1H, d, J = 16 Hz), 7.79 (1H, s); m/z (API): 361.2 (MH<sup>+</sup>; 100%)

5

**Example 3c****E-N-(2-Methyl-1,2,3,4-tetrahydroisoquinolin-7-yl)-3-cinnamide hydrochloride**

The title compound (0.20g) isolated as a pale yellow solid, was prepared from amine D5c (0.16g) and cinnamic acid according to the procedure of Example 1c.

10

<sup>1</sup>H NMR (250MHz, CDCl<sub>3</sub>) δ: 2.43 (3H, s), 2.66 (2H, t), 2.87 (2H, t), 3.52 (2H, s), 6.56 (1H, d), 7.05 (1H, d), 7.26 - 7.52 (6H, m), 7.67(1H, br. s.) and 7.73(1H, d);

MS m/z (API): 293.2 (MH<sup>+</sup>; 100%)

15

**Example 4c****E-N-(2-Methyl-1,2,3,4-tetrahydroisoquinolin-7-yl)-2-methoxycinnamide**

The title compound (0.28g) isolated as a pale yellow solid, was prepared from amine D5c (0.16g) and 2-methoxycinnamic acid (0.18g) according to the procedure of Example 1c

20

<sup>1</sup>H NMR (250MHz, CDCl<sub>3</sub>) δ: 2.39 (3H, s), 2.63 (2H, t), 2.84 (2H, t), 3.51 (2H, s), 3.80 (3H, s), 6.74 (1H, d), 6.85(2H, m), 6.99(1H, d), 7.20 - 7.43(4H, m), 8.00(1H, d) and 8.12(1H, br. s); m/z (API): 323.2 (MH<sup>+</sup>; 100%)

25 **Example 5c****E-N-(2-Methyl-1,2,3,4-tetrahydroisoquinolin-7-yl)-4-chlorocinnamide**

The title compound (0.23g) was prepared from amine D5c (0.16g) and 4-chlorocinnamic acid (0.18g) according to the procedure of Example 1c.

<sup>1</sup>H NMR (250MHz, CDCl<sub>3</sub>) δ: 2.36 (3H, s), 2.60 (2H, t), 2.84 (2H, t), 3.40 (2H, s), 6.62 (1H, d), 6.97 (2H, m), 7.25 (4H, Abq), 7.33 (2H, m), 7.60 (1H, d) and 8.81 (1H, br. s);

30

MS m/z (API): 327, 329 (MH<sup>+</sup>)

**Example 6c**35 **E-N-(2-Methyl-1,2,3,4-tetrahydroisoquinolin-7-yl)-3-chlorocinnamide**

The title compound (0.14g) was prepared from amine D5c (0.16g) and 3-chlorocinnamic acid (0.18g) according to the procedure of Example 1c.

<sup>1</sup>H NMR (250MHz, CDCl<sub>3</sub>) δ: 2.39 (3H, s), 2.62 (2H, t), 2.84 (2H, t), 3.44 (2H, s), 6.62 (1H, d), 7.00 (1H, m), 7.19 - 7.39(6H, m), 7.61 (1H, d) and 8.50 (1H, br. s.);

MS m/z (API): 327, 329 (MH<sup>+</sup>; 100%)

5

**Example 7c****E-N-(2-Methyl-1,2,3,4-tetrahydroisoquinolin-7-yl)-3-methoxycinnamide**

The title compound (0.14g) was prepared from amine D5c (0.16g) and 3-methoxycinnamic acid (0.18g) according to the procedure of Example 1c.

10 <sup>1</sup>H NMR (250MHz, CDCl<sub>3</sub>) δ: 2.39 (3H, s), 2.63 (2H, t), 2.83 (2H, t), 3.43 (2H, s), 3.75 (3H, s), 6.63 (1H, d), 6.85 (1H, d), 6.96 - 7.05 (3H, m), 7.18 - 7.33 (3H, m), 7.67 (1H, d) and 8.41 (1H, br. s); m/z (API): 323 (MH<sup>+</sup>; 100%)

**Example 8c**

15 **E-N-(2-Methyl-1,2,3,4-tetrahydroisoquinolin-7-yl)-α-methylcinnamide**

The title compound (0.14g) was prepared from amine D5c (0.16g) and α-methylcinnamic acid (0.16g) according to the procedure of Example 1c.

<sup>1</sup>H NMR (250MHz, CDCl<sub>3</sub>) δ: 2.14 (3H, s), 2.42 (3H, s), 2.65 (2H, t), 2.87 (2H, t), 3.51 (2H, s), 7.03 (1H, d), 7.26 - 7.41(8H, m) and 7.86 (1H, s); MS m/z (API):  
20 307 (MH<sup>+</sup>; 100%)

**Example 9c****E-N-(2-Methyl-1,2,3,4-tetrahydroisoquinolin-7-yl)-2-chlorocinnamide**

The title compound (0.17g) was prepared from amine D5c (0.16g) and 2-chlorocinnamic acid (0.18g) according to the procedure of Example 1c.

25 <sup>1</sup>H NMR (250MHz, CDCl<sub>3</sub>) δ: 2.31 (3H, s), 2.59 (2H, t), 2.80 (2H, t), 3.40 (2H, s), 6.66 (1H, d), 6.96 (1H, d), 7.09 (1H, t), 7.20 (1H, dt) 7.20 - 7.37 (3H, m), 7.43 (1H, d), 8.09 (1H, d) and 8.83 (1H, br. s.); MS m/z (API): 327, 329 (MH<sup>+</sup>)

30 **Example 10c**

**E-N-(2-Methyl-1,2,3,4-tetrahydroisoquinolin-7-yl)-4-methoxycinnamide**

The title compound (0.18g) was prepared from amine D5c (0.16g) and 4-methoxycinnamic acid (0.18g) according to the procedure of Example 1c.

35 <sup>1</sup>H NMR (250MHz, CDCl<sub>3</sub>) δ: 2.34 (3H, s), 2.58 (2H, t), 2.80 (2H, t), 3.37 (2H, s), 3.73 (3H, s), 6.58 (1H, d), 6.72 (2H, d), 6.94 (1H, m), 7.30 (2H, d), 7.39 (2H, br. s.), 7.65 (1H, d) and 9.00(1H, br. s.); m/z (API): 323 (MH<sup>+</sup>)

**Example 11c****E-N-(6-methyl-5,6,7,8-tetrahydro[1,6]naphthyridin-3-yl)-3-phenylacrylamide**

5 The title compound (0.18g) was prepared from amine D9c (0.16g) and cinnamic acid (0.16g) according to the procedure of Example 1c.

$^1\text{H}$  NMR (250MHz,  $\text{CDCl}_3$ )  $\delta$ : 2.47 (3H, s), 2.77 (2H, t), 3.01 (2H, t), 3.57 (2H, s), 6.46 (1H, d), 7.35 (3H, m), 7.47 (2H, m), 7.73 (1H, d), 8.03 (1H, s) 8.25 (1H, br. s) and 8.40 (1H, s);  $m/z$  (APD): 293 ( $\text{MH}^+$ )

10 **Example 12c**

**E-3-Furan-2-yl-N-(2-methyl-1,2,3,4-tetrahydroisoquinolin-7-yl)acrylamide**

The title compound (0.22g) was prepared from amine D5c (0.16g) and 3-furan-2-yl acrylic acid (0.14g) according to the procedure of Example 1c.

15  $^1\text{H}$  NMR (250MHz,  $\text{CDCl}_3$ )  $\delta$ : 2.39 (3H, s), 2.63 (2H, t), 2.86 (2H, t), 3.42 (2H, s), 6.40 (1H, m), 6.47 (1H, m), 6.54 (1H, d), 6.97 (1H, d), 7.27 - 7.32 (2H, m), 7.41 (1H, s.), 7.48 (1H, d) and 8.48 (1H, br. s.);  $m/z$  (APD): 283 ( $\text{MH}^+$ ; 100%)

**Example 13c****E-N-(2-methyl-1,2,3,4-tetrahydroisoquinolin-7-yl)-3-thiophen-2-ylacrylamide**

20 The title compound (0.23g) was prepared from amine D5c (0.16g) and 3-thiophen-2-yl acrylic acid (0.15g) according to the procedure of Example 1c.

$^1\text{H}$  NMR (250MHz,  $\text{CDCl}_3$ )  $\delta$ : 2.39 (3H, s), 2.63 (2H, t), 2.83 (2H, t), 3.42 (2H, s), 6.45 (1H, d), 6.98 (2H, m), 7.11 (1H, m), 7.27 - 7.30 (3H, m), 7.81 (1H, d) and 8.55 (1H, br. s.).

25 MS  $m/z$  (APD): 299 ( $\text{MH}^+$ ; 100%)

**Example 14c****E-N-(2-Methyl-1,2,3,4-tetrahydroisoquinolin-7-yl)-2, 4-dichlorocinnamide**

30 The title compound (0.36g) was prepared from amine D5c (0.16g) and 2, 4-dichlorocinnamic acid (0.22g) according to the procedure of Example 1c.

$^1\text{H}$  NMR (250MHz,  $d^6$ -DMSO)  $\delta$ : 2.51 (3H, s), 2.89 - 2.96 (4H, m), 3.85 (2H, s), 7.07 (1H, d), 7.12 (1H, d), 7.49 - 7.56 ((3H, m), 7.70 (2H, m), 7.79 (1H, d) and 10.57 (1H, br. s).

MS  $m/z$  (APD): 361, 363 ( $\text{MH}^+$ ; 100%)

35

**Example 15c****Z-N-(2-Methyl-1,2,3,4-tetrahydroisoquinolin-7-yl)-2-methoxycinnamide**

The title compound (0.20g) was prepared from amine D5c (0.16g) and Z-2-methoxycinnamic acid (0.18g) according to the procedure of Example 1c.

<sup>1</sup>H NMR (250MHz, CDCl<sub>3</sub>) δ: 2.42 (3H, s), 2.64 (2H, m), 2.83 (2H, m), 3.49 (2H, s), 3.83 (3H, s), 6.10 (1H, d J = 12Hz), 6.90 - 7.32 (8H, m) and 7.44(1H, d);

5 MS m/z (API): 323 (MH<sup>+</sup>; 100%)

#### Example 16c

##### E-3-Indolin-5-yl-N-(2-Methyl-1,2,3,4-tetrahydroisoquinolin-7-yl)-acrylamide

10 The title compound (0.14g) was prepared from amine D5c (0.16g) and E-3-indolin-5-yl acrylic acid (0.19g) according to the procedure of Example 1c.

<sup>1</sup>H NMR (250MHz, d<sup>6</sup>-DMSO) δ: 2.70 (3H, s), 2.95 (2H, m), 3.16 (2H, m), 4.04 (2H, s), 6.50 (1H, s), 6.75 (1H, d), 7.15 (1H, d), 7.41 - 7.57 (5H, m), 7.65 (1H, d), 7.79 (1H, s), 10.18 (1H, s) and 11.35 (1H, s); m/z (API): 332 (MH<sup>+</sup>)

#### 15 Example 17c

##### E-3-(1-Methyl-Indolin-2-yl)-N-(2-Methyl-1,2,3,4-tetrahydroisoquinolin-7-yl)-acrylamide

20 The title compound (0.20g) was prepared from amine D5c (0.16g) and E-3-(1-methyl-indolin-2-yl) acrylic acid (0.20g) according to the procedure of Example 1c.

<sup>1</sup>H NMR (250MHz, CDCl<sub>3</sub>) δ: 2.43 (3H, s), 2.66 (2H, t), 2.88 (2H, t), 3.54 (2H, s), 3.77 (3H, s), 6.60 (1H, d), 6.89 (1H, s), 7.05 (2H, m), 7.21 - 7.31 (2H, m), 7.42 (1H, br.s), 7.56 (1H, d), 7.62 (1H, br. s) and 7.86 (1H, d); m/z (API): 345 (MH<sup>+</sup>; 100%)

25

#### Example 18c

##### E-N-(2-Methyl-1,2,3,4-tetrahydroisoquinolin-7-yl)-3-chloro-4-methoxycinnamide

30 The title compound (0.37g) was prepared from amine D5c (0.16g) and E-3-chloro-4-methoxycinnamic acid (0.22g) according to the procedure of Example 1c.

<sup>1</sup>H NMR (250MHz, d<sup>6</sup>-DMSO) δ: 2.87 (3H, s), 3.05 (2H, m), 3.38 (2H, m), 3.91 (3H, s), 4.34 (2H, s), 6.81 (1H, d), 7.21 (2H, m), 7.47 - 7.71 (5H, m) and 10.35 (1H, br. s);

MS m/z (API): 357, 359 (MH<sup>+</sup>; 100%)

35

#### Example 19c

**E-N-(2-Methyl-1,2,3,4-tetrahydroisoquinolin-7-yl)-4-methylsulphonylcinnamide**

The title compound (0.17g) was prepared from amine D5c (0.16g) and E-4-methylsulphonylcinnamic acid (0.22g) according to the procedure of Example 1c.

- 5  $^1\text{H}$  NMR (250MHz,  $\text{d}^6$ -DMSO)  $\delta$ : 2.44 (3H, s), 2.74 - 2.82 (4H, m), 3.26 (3H, s), 3.63 (2H, s), 7.01 (1H, d), 7.10 (1H, d), 7.47 (2H, m), 7.64 (1H, d), 7.87 (2H, d), 7.98 (2H, d) and 10.32 (1H, br. s);  $m/z$  (API): 371 ( $\text{MH}^+$ ; 100%)

**Example 20c**

- 10 **E-N-methyl-3-[2-(2-methyl-1,2,3,4-tetrahydroisoquinolin-7-ylcarbamoyl)vinyl] benzamide**

The title compound (0.17g) was prepared from amine D5c (0.16g) and E-3-(3-methylcarbamoylphenyl)acrylic acid (0.21g) according to the procedure of Example 1c.

- 15 MS  $m/z$  (API): 349 ( $\text{MH}^+$ ; 100%)

**Example 21c****E-3-(Indazolin-3-yl)-N-(2-Methyl-1,2,3,4-tetrahydroisoquinolin-7-yl)-acrylamide**

- 20 The title compound (0.05) was prepared from amine D5c (0.16g) and E-3-indazolin-3-yl acrylic acid (0.19g) according to the procedure of Example 1c.

- $^1\text{H}$  NMR (250MHz,  $\text{d}^6$ -DMSO)  $\delta$ : 2.38 (3H, s), 2.65 (2H, m), 2.79 (2H, m), 3.54 (2H, s), 7.09 (1H, d), 7.22 (1H, d), 7.30 (1H, m), 7.47 (2H, m), 7.62 (1H, d), 7.79 (1H, d), 8.09 (1H, d), 10.16 (1H, br. s) and 13.52 (1H, br. s);  $m/z$  (API): 332 (MH<sup>+</sup>; 100%)

**Example 22c****E-N-(2-Methyl-1,2,3,4-tetrahydroisoquinolin-7-yl)-2-methylcinnamide**

- 30 The title compound (0.08g) was prepared from amine D5c (0.16g) and E-2-methylcinnamic acid (0.16g) according to the procedure of Example 1c.

- $^1\text{H}$  NMR (250MHz,  $\text{d}^6$ -DMSO)  $\delta$ : 2.24 (3H, s), 2.32 (3H, s), 2.48 (2H, t), 2.67 (2H, m), 6.64 (1H, d), 6.97 (1H, d), 7.20 (3H, m), 7.33 (2H, m), 7.49 (1H, d), 7.70 (1H, d) and 10.03 (1H, br. s);  $m/z$  (API): 307 ( $\text{MH}^+$ ; 100%)

**Example 23c****E-N-(2-Methyl-1,2,3,4-tetrahydroisoquinolin-7-yl)-2-nitrocinnamide**

- The title compound (0.04g) was prepared from amine D5c (0.16g) and E-2-nitrocinnamic acid (0.19g) according to the procedure of Example 1c.

<sup>1</sup>H NMR (250MHz, CDCl<sub>3</sub>) δ: 2.46 (3H, s), 2.68 (2H, s), 2.90 (2H, t), 3.57 (2H, s), 6.45 (1H, d), 7.08 (1H, d), 7.53 (2H, br. s), 7.53 (1H, m), 7.63 (2H, s), 8.04 (1H, d) and 8.10 (1H, d); m/z (APD): 338 (MH<sup>+</sup>; 100%)

5 **Example 24c**

**E-N-(2-Methyl-1,2,3,4-tetrahydroisoquinolin-7-yl)-2-trifluoromethylcinnamide**

The title compound (0.05g) was prepared from amine D5c (0.16g) and E-2-trifluoromethylcinnamic acid (0.22g) according to the procedure of Example 1c.

10 <sup>1</sup>H NMR (250MHz, CDCl<sub>3</sub>) δ: 2.37 (3H, s), 2.60 (2H, s), 2.80 (2H, m), 3.41 (2H, s), 6.59 (1H, d), 6.95 (1H, d), 7.26 (1H, d), 7.35 - 7.40 (3H, m), 7.54 (1H, d), 7.64 (1H, d), 8.08 (1H, d) and 8.69 (1H, br. s); m/z (APD): 361 (MH<sup>+</sup>; 100%)

**Example 25c**

15 **E-N-(2-Methyl-1,2,3,4-tetrahydroisoquinolin-7-yl)-2-ethoxycinnamide**

The title compound (0.06g) was prepared from amine D5c (0.16g) and E-2-ethoxycinnamic acid (0.19g) according to the procedure of Example 1c.

20 <sup>1</sup>H NMR (250MHz, CDCl<sub>3</sub>) δ: 1.41 (3H, t), 2.63 (2H, t), 2.84 (2H, m), 3.46 (2H, s), 4.40 (2H, q), 6.70 (1H, d), 6.82 - 6.87 (2H, m), 6.99 (1H, d), 7.23 - 7.45 (4H, m), 8.05 (1H, d) and 8.13 (1H, br. s); m/z (APD): 337 (MH<sup>+</sup>; 100%)

**Example 26c**

**E-N-(2-Methyl-1,2,3,4-tetrahydroisoquinolin-7-yl)-2-chloro-4-fluorocinnamide**

25 The title compound (0.13g) was prepared from amine D5c (0.16g) and E-2-chloro-4-fluorocinnamic acid (0.20g) according to the procedure of Example 1c.

<sup>1</sup>H NMR (250MHz, CDCl<sub>3</sub>) δ: 2.46 (3H, s), 2.69 (2H, t), 2.89 (2H, m), 3.57 (2H, s), 6.48 (1H, d), 6.99 (1H, dt), 7.08 (1H, d), 7.18 (1H, dd), 7.26 (1H, m), 7.35 (1H, s), 7.43 (1H, s), 7.58 (1H, m) and 8.04 (1H, d); m/z (APD): 345 (MH<sup>+</sup>; 100%)

30

**Example 27c**

**E-N-(2-Methyl-1,2,3,4-tetrahydroisoquinolin-7-yl)-2-chloro-6-fluorocinnamide**

The title compound (0.06g) was prepared from amine D5c (0.16g) and E-2-chloro-6-fluorocinnamic acid (0.20g) according to the procedure of Example 1c.

35

<sup>1</sup>H NMR (250MHz, CDCl<sub>3</sub>) δ: 2.44 (3H, s), 2.67 (2H, t), 2.88 (2H, m), 3.54 (2H, s), 6.83 (1H, d), 6.99 - 7.07 (2H, m), 7.18 - 7.29 (4H, m), 7.41 (1H, s), 7.70(s, 1H and 7.96 (1H, d); m/z (API): 345 (MH<sup>+</sup>; 100%)

5 **Example 28c**

**E-N-(2-Methyl-1,2,3,4-tetrahydroisoquinolin-5-yl)-4-chlorocinnamide**

The title compound (0.02g) was prepared from amine **D11c** (0.16g) and E-4-chlorocinnamic acid (0.18g) according to the procedure of Example 1c.

10 <sup>1</sup>H NMR (250MHz, CDCl<sub>3</sub>) δ: 2.47 (3H, s), 2.74 - 2.79 (4H, m), 3.87 (2H, s), 6.52 (1H, d), 6.91 (1H, d), 7.03 (1H, s), 7.19 (1H, t), 7.35 (2H, d), 7.45 (2H, d), 7.70 (1H, d) and 7.79 (1H, br. s); m/z (API): 327, 329 (MH<sup>+</sup>; 100%)

**Example 29c**

**E-N-(2-Methyl-1,2,3,4-tetrahydroisoquinolin-5-yl)-cinnamide**

15 The title compound (0.14g) was prepared from amine **D11c** (0.16g) and cinnamic acid (0.15g) according to the procedure of Example 1c.

<sup>1</sup>H NMR (250MHz, CDCl<sub>3</sub>) δ: 2.35 (3H, s), 2.62(2H, t), 2.73 (2H, t), 3.52 (2H, s), 6.66 (1H, d), 6.81 (1H, d), 7.07 (1H, t), 7.31 (3H, m), 7.44 (3H, m), 7.66 (1H, d) and 7.98 (1H, br. s); m/z (API): 293 (MH<sup>+</sup>; 100%)

20

**Example 30c**

**E-N-(2-Methyl-1,2,3,4-tetrahydroisoquinolin-5-yl)-3-chlorocinnamide**

The title compound (0.06) was prepared from amine **D11c** (0.16g) and 3-chlorocinnamic acid (0.18g) according to the procedure of Example 1c.

25 <sup>1</sup>H NMR (250MHz, CDCl<sub>3</sub>) δ: 2.49 (3H, s), 2.80 (4H, m), 3.64 (2H, s), 6.57 (1H, d), 6.91 (1H, d), 7.19 (1H, t), 7.36 (4H, m), 7.51 (1H, br. s.), 7.68 (1H, d) and 7.79 (1H, br. s);

MS m/z (API): 327, 329 (MH<sup>+</sup>; 100%)

30 **Example 31c**

**E-N-(2-Methyl-1,2,3,4-tetrahydroisoquinolin-5-yl)-2-chlorocinnamide**

The title compound (0.10) was prepared from amine **D11c** (0.16g) and 2-chlorocinnamic acid (0.18g) according to the procedure of Example 1c.

35 <sup>1</sup>H NMR (250MHz, CDCl<sub>3</sub>) δ: 2.47 (3H, s), 2.75 (4H, m), 3.60 (2H, s), 6.57 (1H, d), 6.91 (1H, d), 7.10 (1H, br. s), 7.19 (1H, t), 7.30 (2H, m), 7.41 (1H, m), 7.62 (1H, br. s), 7.90 (1H, br. s) and 8.13 (1H, d); m/z (API): 327, 329 (MH<sup>+</sup>; 100%)

**Example 32c****E-N-(2-Methyl-1,2,3,4-tetrahydroisoquinolin-5-yl)-3-acetylcinnamide hydrochloride**

5 The title compound (0.12) was prepared from amine **D11c** (0.16g) and 3-acetylcinnamic acid (0.19g) according to the procedure of Example 1c. The hydrochloride salt was prepared by treatment of the free base in methanol with diethyl ether/HCl.

<sup>1</sup>H NMR (250MHz, CDCl<sub>3</sub>) δ: 2.47 (3H, s), 2.64 (3H, s), 2.74 - 2.81 (4H, m), 3.60 (2H, s), 6.69 (1H, d), 6.90 (1H, d), 7.19 (2H, m), 7.49 (1H, t), 7.69 (1H, m),  
10 7.78 (1H, d), 7.93 (1H, d) and 8.15 (1H, br. s); m/z (APD): 335 (MH<sup>+</sup>; 100%)

**Example 33c****E-N-(2-Methyl-1,2,3,4-tetrahydroisoquinolin-5-yl)-2-bromocinnamide hydrochloride**

15 The title compound (0.10) was prepared from amine **D11c** (0.16g) and 2-bromocinnamic acid (0.23g) according to the procedure of Example 1c. The hydrochloride salt was prepared by treatment of the free base in methanol with diethyl ether/HCl.

<sup>1</sup>H NMR (250MHz, CDCl<sub>3</sub>) δ: 2.46 (3H, s), 2.71 - 2.80 (4H, m), 3.59 (2H, s),  
20 6.51 (1H, d), 6.91 (1H, d), 7.07 (1H, br. s.), 7.16 - 7.26 (2H, m), 7.32 (1H, t), 7.60 (1H, br. s.), 7.61 (1H, d), 7.86 (1H, br. s); m/z (APD): 371, 373 (MH<sup>+</sup>; 100%)

**Example 34c****E-N-(2-Methyl-1,2,3,4-tetrahydroisoquinolin-5-yl)-2-methylcinnamide hydrochloride**

25 The title compound (0.05) was prepared from amine **D11c** (0.16g) and 2-methylcinnamic acid (0.16g) according to the procedure of Example 1c. The hydrochloride salt was prepared by treatment of the free base in methanol with diethyl ether/HCl.

30 <sup>1</sup>H NMR (250MHz, CDCl<sub>3</sub>) δ: 2.46 (3H, s), 2.75 (3H, s), 3.03 (2H, m), 3.13 (2H, m), 3.99 (2H, s), 6.54 (1H, d), 6.89 (1H, d), 7.18 - 7.42 (6H, m), 7.59 (1H, d), 8.04 (1H, d);  
MS m/z (APD): 307 (MH<sup>+</sup>; 100%)

**Example 35 c****E-N-(2-Methyl-1,2,3,4-tetrahydroisoquinolin-5-yl)-4-ethoxycinnamide**

The title compound (0.08g) was prepared from amine **D11c** (0.16g) and 4-ethoxycinnamic acid (0.19g) according to the procedure of Example 1c.



<sup>1</sup>H NMR (250MHz, CDCl<sub>3</sub>) δ: 1.43 (3H, t), 2.47 (3H, s), 2.74 - 2.80 (4H, m), 3.60 (2H, s), 4.06 (2H, q), 6.42 (1H, d), 6.87 - 6.96 (4H, m), 7.19 (1H, t), 7.47 (2H, d), 7.71 (1H, d) and 7.82 (1H, br. s.); m/z (APD): 337 (MH<sup>+</sup>; 100%)

5    **Example 36c**

**E-N-(2-Methyl-1,2,3,4-tetrahydroisoquinolin-5-yl)-2-methoxycinnamide**

The title compound (0.07g) was prepared from amine D11c (0.16g) and 2-methoxycinnamic acid (0.18g) according to the procedure of Example 1c.

10    <sup>1</sup>H NMR (250MHz, CDCl<sub>3</sub>) δ: 2.47 (3H, s), 2.74 - 2.80 (4H, m), 3.60 (2H, s), 3.90 (3H, s), 6.69 (1H, d), 6.88 - 7.02 (4H, m), 7.18 (1H, t), 7.34 (1H, t), 7.50 (1H, d), 7.86 (1H, br. s) and 8.01 (1H, d); m/z (APD): 323 (MH<sup>+</sup>; 100%)

**Example 37c**

15    **E-N-(2-Methyl-1,2,3,4-tetrahydroisoquinolin-5-yl)-5-bromo-2-methoxycinnamide**

The title compound (0.16g) was prepared from amine D11c (0.16g) and 5-bromo-2-methoxycinnamic acid (0.26g) according to the procedure of Example 1c.

20    <sup>1</sup>H NMR (250MHz, CDCl<sub>3</sub>) δ: 2.47 (3H, s), 2.71 - 2.81 (4H, m), 3.60 (2H, s), 3.87 (3H, s), 6.62 (1H, d), 6.80 (1H, d), 6.90 (1H, d), 7.00 (1H, br. s), 7.19 (1H, t), 7.41 (1H, dd), 7.61 (1H, br. s), 7.85 (1H, br. s) and 7.95 (1H, d); m/z (APD): 401, 403 (MH<sup>+</sup>; 100%)

**Example 38c**

25    **E-2-Cyano-N-(2-Methyl-1,2,3,4-tetrahydroisoquinolin-5-yl)cinnamide hydrochloride**

The title compound (0.11) was prepared from amine D11c (0.16g) and 4-bromo-2-cyanocinnamic acid (0.19g) according to the procedure of Example 1c. The hydrochloride was prepared from the free base in MeOH methanol and diethyl ether/HCl.

30    <sup>1</sup>H NMR (250MHz, CDCl<sub>3</sub>) δ: 2.47 (3H, s), 2.74 - 2.80 (4H, m), 3.60 (2H, s), 6.82 - 6.93 (2H, m), 7.16 - 7.22 (2H, m), 7.46 (1H, t), 7.59 - 7.73 (3H, m), 7.83 (1H, br. s.), 7.96 (1H, d); m/z (APD): 318 (MH<sup>+</sup>; 100%)

**Example 39c**

35    **N-(8-Chloro-2-trifluoroacetyl-1,2,3,4-tetrahydroisoquinolin-7-yl)-2-chlorocinnamide**

The title compound (0.13g) was prepared from amine **D13c** (0.21g) and 2-chlorocinnamoyl chloride (0.45g) according to the procedure of Example 2c.

<sup>1</sup>H NMR (250MHz, CDCl<sub>3</sub>) δ: 2.97 (2H, m), 3.82 - 3.93 (2H, m), 4.80 (2H, d due to restricted amide rotation), 6.60 (1H, d), 7.16 (1H, t), 7.33 (2H, m), 7.46 (1H, m),  
5 7.67 (1H, m), 7.84 (1H, d), 8.18 (1H, d) and 8.43 (1H, d); m/z (API): 443, 445 (MH<sup>+</sup>; 100%)

#### Example 40c

##### N-(8-Chloro-1,2,3,4-tetrahydroisoquinolin-7-yl)-2-chlorocinnamide

10 A solution of N-(8-Chloro-2-trifluoroacetyl-1,2,3,4-tetrahydroisoquinolin-7-yl)-2-chlorocinnamide (0.2gg) in methanol/water (5ml 9:1) was treated with potassium carbonate (0.38g) and stirred 12h. The mixture was diluted with dichloromethane and washed with water. The organic phase was dried (MgSO<sub>4</sub>), solvent removed at reduced pressure. The residue was column chromatographed (silica gel,  
15 dichloromethane/methanol/ammonia upto 9:1:0.1 eluant) to give the title compound (0.10g) as a colourless solid. <sup>1</sup>H NMR (250MHz, CDCl<sub>3</sub>) δ: 2.78 (2H, t), 3.10 (2H, t), 4.03 (2H, s), 6.59 (1H, d), 7.07 (1H, d), 7.28 - 7.32 (2H, m), 7.42 (1H, m), 7.66 (1H, m), 7.82 (1H, br. s), 8.16 (1H, d) and 8.31 (1H, d);  
m/z (API): 347, 349 (MH<sup>+</sup>; 100%)

20

#### Example 41c

##### N-(8-Chloro-2-methyl-1,2,3,4-tetrahydroisoquinolin-7-yl)-2-chlorocinnamide

A solution of N-(8-Chloro-2-methyl-1,2,3,4-tetrahydroisoquinolin-7-yl)-2-chlorocinnamide (0.08g) in 37% aqueous formaldehyde (0.63ml) and formic acid  
25 (0.34ml) and stirred at 80°C for 3h. Solid sodium hydroxide was added to neutralise the solution and the aqueous phase extracted with dichloromethane. The combined organic extracts were dried (MgSO<sub>4</sub>) and solvent removed at reduced pressure to give the title compound (0.07g).

<sup>1</sup>H NMR (250MHz, CDCl<sub>3</sub>) δ: 2.51 (3H, s), 2.66 (2H, t), 2.90 (2H, t), 3.59 (2H, s), 6.59 (1H, d), 7.07 (1H, d), 7.27 - 7.31 (2H, m), 7.41 (1H, m), 7.64 (1H, m),  
30 7.86 (1H, br. s), 8.14 (1H, d) and 8.27 (1H, d); m/z (API): 361, 363 (MH<sup>+</sup>; 100%)

#### Example 42c

##### N-(8-Bromo-2-trifluoroacetyl-1,2,3,4-tetrahydroisoquinolin-7-yl)-2-chlorocinnamide

35 The title compound (0.27g) was prepared from amine **D14c** (0.32g) and 2-chlorocinnamoyl chloride according to the procedure of Example 2c.

<sup>1</sup>H NMR (250MHz, CDCl<sub>3</sub>) δ: 2.93 - 3.00 (2H, m), 3.82 - 3.92 (2H, m due to restricted rotation), 4.77 (2H, d, due to restricted rotation) 6.60 (1H, d), 7.17 - 7.36 (3H, m), 7.40 - 7.47 (1H, m), 7.40 - 7.47 (1H, m), 7.87 (1H, m), 8.18 (1H, d) and 8.39 (1H, d);

5 MS m/z (API): 361, 363 (MH<sup>+</sup>; 100%)

#### Example 43c

##### E-N-(2-Methyl-1,2,3,4-tetrahydroisoquinolin-7-yl)-α-fluorocinnamide

The title compound (0.20g) was prepared from amine D5c (0.17g) and Z-α-fluorocinnamic acid (0.19g) according to the procedure of Example 1c. <sup>1</sup>H NMR (250MHz, CDCl<sub>3</sub>) δ: 2.44 (3H, s), 2.67 (2H, t), 2.89 (2H, t), 3.55 (2H, s), 7.03 (1H, d, J = 39Hz), 7.08 (1H, d), 7.29 - 7.41 (4H, m), 7.63 (2H, m) and 8.16 (1H, d); m/z (API): 311 (MH<sup>+</sup>; 100%)

15 The following Examples were made in a manner similar to the procedures described in the above Descriptions and Examples

#### Example 44c

##### E-N-(8-Bromo-1,2,3,4-tetrahydroisoquinolin-7-yl)-2-chlorocinnamide

20 The title compound (0.36g) was prepared from the trifluoroacetamide of Example 42 (0.681g) according to the method of Example 40.

MS m/z (API<sup>+</sup>): 391, 393. (MH)<sup>+</sup>

#### Example 45c

25 E-N-(8-Bromo-2-methyl-1,2,3,4-tetrahydroisoquinolin-7-yl)-2-chlorocinnamide

The title compound (0.28g) was prepared from the amine of Example 44c (0.361g) according to the method of Example 41c.

MS m/z (API<sup>+</sup>): 405, 407 (MH)<sup>+</sup>

30

#### Example 46c

##### E-N-(2,4,4-Trimethyl-1,2,3,4-tetrahydroisoquinolin-7-yl)-2-chlorocinnamide

<sup>1</sup>H NMR (CDCl<sub>3</sub>) δ: 1.30 (6H, s), 2.39 (2H, s), 2.41 (3H, s), 3.53 (2H, s), 6.53 (1H, d), 7.40 (7H, m), 7.59 (1H, m), 8.11 (1H, d); m/z: (API<sup>+</sup>): 355.2 (MH<sup>+</sup>;

35 100%)

#### Example 47c

**E-N-(8-Ethyl-2,4,4-trimethyl-1,2,3,4-tetrahydroisoquinolin-7-yl)-2-methylcinnamide**

<sup>1</sup>H NMR (CDCl<sub>3</sub>) δ: 1.15 (3H, t), 1.35 (6H, s), 2.50 (10H, m), 3.66 (2H, s), 6.49 (1H, d), 7.21 (6H, m), 7.55 (1H, s), 8.05 (1H, d); m/z (AP<sup>+</sup>): 363.3 (MH<sup>+</sup>; 100%)

5

**Example 48c**

**E-N-(8-Ethyl-2,4,4-trimethyl-1,2,3,4-tetrahydroisoquinolin-7-yl)-2-chloro-6-fluorocinnamide**

<sup>1</sup>H NMR (CDCl<sub>3</sub>) δ: 1.15 (3H, t), 1.32 (6H, s), 2.40 (2H, s), 2.48 (3H, s), 2.59 (2H, q), 3.56 (2H, s), 6.87 (1H, d), 7.21 (5H, m), 7.67 (1H, s), 7.98 (1H, s); m/z (AP<sup>+</sup>): 401.2 (MH<sup>+</sup>; 100%)

10

**Example 49c**

**E-N-(8-Ethyl-2,4,4-trimethyl-1,2,3,4-tetrahydroisoquinolin-7-yl)-2-trifluoromethylcinnamide**

<sup>1</sup>H NMR (CDCl<sub>3</sub>) δ: 1.15 (3H, t), 1.34 (6H, s), 2.54 (7H, m), 3.66 (2H, s), 6.59 (1H, d), 7.23 (1H, d), 7.43 (3H, m), 7.69 (2H, t), 8.06 (1H, d); m/z (AP<sup>+</sup>): 417.2 (MH<sup>+</sup>; 100%).

15

**Example 50c**

**E-N-(8-Ethyl-2,4,4-trimethyl-1,2,3,4-tetrahydroisoquinolin-7-yl)-2-chloro-4-fluorocinnamide**

<sup>1</sup>H NMR (CDCl<sub>3</sub>) δ: 1.13 (3H, t), 1.34 (6H, s), 2.53 (7H, m), 3.64 (2H, s), 6.60 (1H, d), 6.99 (2H, m), 7.18 (2H, m), 7.44 (1H, s), 7.60 (2H, m), 8.04 (1H, d); m/z (AP<sup>+</sup>): 401.2 (MH<sup>+</sup>; 100%).

20

25

**Example 51c**

**E-N-(8-Ethyl-2,4,4-trimethyl-1,2,3,4-tetrahydroisoquinolin-7-yl)-2-chlorocinnamide**

<sup>1</sup>H NMR (CDCl<sub>3</sub>) δ: 1.16 (3H, t), 1.33 (6H, s), 2.46 (2H, s), 2.58 (5H, m), 3.62 (2H, s), 6.64 (1H, d), 7.30 (6H, m), 7.63 (1H, s), 8.11 (1H, d); m/z (AP<sup>+</sup>): 383.1 (MH<sup>+</sup>; 100%)

30

**Example 52c**

**E-N-(1,1,2-Trimethyl-1,2,3,4-tetrahydroisoquinolin-7-yl)-2-chlorocinnamide**

<sup>1</sup>H NMR (CDCl<sub>3</sub>) δ: 1.44 (6H, s), 2.47 (3H, s), 2.81 - 2.95 (4H, m), 6.62 (1H, d, J = 16 Hz), 6.95 - 7.07 (1H, m), 7.20 - 7.44 (4H, m), 7.55 - 7.63 (1H, m), 7.71 - 7.91 (2H, m), 8.12 (1H, d, J = 16 Hz); m/z (AP<sup>+</sup>): 355, 357 (MH<sup>+</sup>)

35

**Example 53c****E-N-(1,2,4,4-Tetramethyl-1,2,3,4-tetrahydroisoquinolin-7-yl)-2-chlorocinnamide**

- 5  $^1\text{H}$  NMR ( $\text{CDCl}_3$ )  $\delta$ : 1.25 (3H, s), 1.28 (3H, s), 1.33 (3H, d,  $J = 7$  Hz), 2.30 (1H, d,  $J = 12$  Hz), 2.43 (3H, s), 2.60 (1H, d,  $J = 12$  Hz), 3.48 (1H, q,  $J = 7$  Hz), 6.59 (1H, d,  $J = 16$  Hz), 7.15 - 7.50 (6H, m), 7.56 (1H, dd,  $J = 8, 2$  Hz), 7.72 (1H, brs), 8.11 (1H, d,  $J = 16$  Hz);  $m/z$  ( $\text{API}^+$ ): 369, 371 ( $\text{MH}^+$ ).

**Example 54c**

- 10 **E-N-(8-Ethyl-2,4,4-trimethyl-1,2,3,4-tetrahydroisoquinolin-7-yl)-2-ethoxycinnamide**

$^1\text{H}$  NMR ( $\text{CDCl}_3$ )  $\delta$ : 1.09 (3H, t), 1.34 (6H, d), 1.48 (3H, t), 2.40 (2H, s), 2.46 (5H, m), 3.57 (2H, s), 4.02 (2H, q), 6.89 (4H, m), 7.15 (2H, m), 7.30 (1H, m), 7.45 (1H, dd), 8.08 (1H, d);  $m/z$  ( $\text{API}^+$ ): 393.3 ( $\text{MH}^+$ ; 50%)

15

**Example 55c****E-N-(8-Chloro-2,4,4-trimethyl-1,2,3,4-tetrahydroisoquinolin-7-yl)-2-chlorocinnamide**

- 20  $^1\text{H}$  NMR ( $\text{CDCl}_3$ )  $\delta$ : 1.30 (6H, s), 2.38 (2H, s), 2.47 (3H, s), 3.57 (2H, s), 6.59 (1H, d), 7.29 (3H, m), 7.43 (1H, m), 7.66 (1H, m), 7.81 (1H, s), 8.15 (1H, d), 8.35 (1H, d);  $m/z$  ( $\text{API}^+$ ): 389.0 ( $\text{M}^+$ ; 95%)

**Example 56c**

- 25 **E-N-(8-Ethyl-2,4,4-trimethyl-1,2,3,4-tetrahydroisoquinolin-7-yl)-3-chloro-4-methoxycinnamide**

$^1\text{H}$  NMR ( $\text{CDCl}_3$ )  $\delta$ : 1.13 (3H, t), 1.34 (6H, s), 2.53 (7H, m), 3.64 (2H, s), 3.92 (1H, s), 6.34 (1H, d), 6.51 (1H, d), 6.91 (1H, d), 7.23 (1H, d), 7.36 (2H, m), 7.61 (2H, m);  $m/z$  ( $\text{API}^+$ ): 413.2 ( $\text{MH}^+$ ; 100%).

- 30 **Example 57c**

**E-N-(8-Ethyl-2,4,4-trimethyl-1,2,3,4-tetrahydroisoquinolin-7-yl)-2-cyanocinnamide**

- 35  $^1\text{H}$  NMR ( $\text{CDCl}_3$ )  $\delta$ : 1.16 (3H, t), 1.31 (6H, s), 2.38 (2H, s), 2.47 (3H, s), 2.57 (2H, q), 3.55 (2H, s), 6.88 (1H, d), 7.24 (1H, d), 7.46 (1H, t), 7.68 (5H, m), 7.95 (1H, d);  $m/z$  ( $\text{API}^+$ ): 374.2 ( $\text{MH}^+$ ; 100%).

**Example 58c**

**E-N-(8-Methyl-2,4,4-trimethyl-1,2,3,4-tetrahydroisoquinolin-7-yl)-2-chlorocinnamide**

<sup>1</sup>H NMR (CDCl<sub>3</sub>) δ: 1.31 (6H, s), 2.11 (3H, s), 2.38 (2H, s), 2.47 (3H, s), 3.47 (2H, s), 6.61 (1H, d), 7.24 (3H, m), 7.52 (4H, m), 8.11 (1H, d);

5 MS m/z (API<sup>+</sup>): 369.2 (MH<sup>+</sup>; 100%).

**Example 59c****E-N-(8-Ethyl-2,4,4-trimethyl-1,2,3,4-tetrahydroisoquinolin-7-yl)-2-acetylcinnamide**

10 <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ: 1.18 (3H, t), 1.31 (6H, s), 2.40 (3H, s), 2.47 (3H, s), 2.61 (5H, m), 3.57 (2H, s), 6.29 (1H, d), 7.23 (1H, d), 7.45 - 7.72 (6H, m), 8.09 (1H, d);  
MS m/z (API<sup>+</sup>): 391.2 (MH<sup>+</sup>; 100%).

**Example 60c**

15 **E-N-(8-Ethyl-4,4-dimethyl-1,2,3,4-tetrahydroisoquinolin-7-yl)-2-chlorocinnamide**

<sup>1</sup>H NMR (CDCl<sub>3</sub>) δ: 1.14 (3H, t), 1.28 (6H, s), 2.56 (2H, q), 2.83 (2H, s), 4.04 (2H, s), 6.60 (1H, d), 7.27 (5H, m), 7.41 (1H, m), 7.64 (1H, s), 8.11 (1H, d);  
MS m/z (API<sup>+</sup>): 369.3 (MH<sup>+</sup>; 100%)

20

**Example 61c****E-N-(8-Ethyl-4,4-dimethyl-1,2,3,4-tetrahydroisoquinolin-7-yl)-2-cyanocinnamide**

25 <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ: 1.13 (3H, t), 1.29 (6H, s), 2.55 (2H, q), 2.82 (2H, s), 4.03 (2H, s), 6.89 (1H, d), 7.24 (1H, d), 7.45 (2H, m), 7.61 (2H, m), 7.71 (1H, d), 7.94 (1H, d);  
MS m/z (API<sup>+</sup>): 360.2 (MH<sup>+</sup>; 100%).

**Example 62c**

30 **E-N-(8-Chloro-2,3,3-trimethyl-1,2,3,4-tetrahydroisoquinolin-7-yl)-2-chlorocinnamide**

<sup>1</sup>H NMR (CDCl<sub>3</sub>) δ: 1.09 (6H, s), 2.42 (3H, s), 2.71 (2H, s), 3.72 (2H, s), 6.59 (1H, d, J = 16 Hz), 7.03 (1H, d, J = 8 Hz), 7.23 - 7.40 (2H, m), 7.44 (1H, m), 7.67 (1H, m), 7.82 (1H, brs), 8.15 (1H, d, J = 16 Hz), 8.30 (1H, brd, J = 8 Hz); m/z  
35 (API<sup>+</sup>): 389.3 (MH<sup>+</sup>; 100%)

**Example 63c**

**E-N-(5-Bromo-1,2,3,4-tetrahydroisoquinolin-7-yl)-2-chlorocinnamide**

MS  $m/z$  (API<sup>+</sup>): 392.9 (MH<sup>+</sup>; 100%, C<sub>18</sub>H<sub>16</sub>BrClN<sub>2</sub> O requires M<sup>+</sup> 391).

**Example 64c**5 **E-N-(5,6,7,8-Tetrahydro-6-methyl[1,6]naphthyridin-3-yl)-cinnamide**

Prepared from *trans*-cinnamic acid and D9c

MS  $m/z$  (API<sup>+</sup>): 294 (MH<sup>+</sup>; 100%, C<sub>18</sub>H<sub>19</sub>N<sub>3</sub> O requires M<sup>+</sup> 293).

**Example 65c**10 **E-N-(5,6,7,8-Tetrahydro-6,8,8-trimethyl[1,6]naphthyridin-3-yl)-2-chlorocinnamide hydrochloride**

Prepared from D47c and *trans*-2-chlorocinnamic acid (183mg; 1.0 mmol) and isolated as a white powder (86mg; 22%).

<sup>1</sup>H NMR [free base] (250 MHz; CD<sub>3</sub>OD)  $\delta$ : 1.24 (6H, s), 2.36 (3H, s), 2.50 (2H, s) 3.50 (2H, s), 6.68 (1H, d, J = 16 Hz), 7.17 – 7.37 (3H, m), 7.64 (1H, m), 7.83 (1H, d, J = 2 Hz), 7.98 (1H, d, J = 16 Hz), 8.50 (1H, d, J = 2 Hz);  
 15  $m/z$  (API<sup>+</sup>): 356.1 (MH)<sup>+</sup>, 378.1 (M+Na)<sup>+</sup>.

**Description 1rc**20 **E-7-(2-Ethoxycarbonylvinyl)-3,4-dihydro-1H-isoquinoline-2-carboxylic acid *tert* butyl ester**

(a) A mixture of 7-bromo-3,4-dihydro-1H-isoquinoline-2-carboxylic acid *tert* butyl ester (1.0g), palladium (II) acetate 0.037g, tris(o-tolyl)phosphine (0.1g) triethylamine (0.67ml) and ethyl acrylate (0.52ml) in acetonitrile (2ml) was boiled  
 25 for 4 h. After cooling to room temperature the mixture was diluted with ethyl acetate, washed with water and brine dried (MgSO<sub>4</sub>) and solvent removed at reduced pressure. The residue was column chromatographed (silica gel, ethyl acetate/hexane) to give after combining of appropriate fractions 7-(2-ethoxycarbonylvinyl)-3,4-dihydro-1H-isoquinoline-2-carboxylic acid *tert* butyl  
 30 ester (0.51g).

<sup>1</sup>H NMR (250MHz CDCl<sub>3</sub>)  $\delta$ : 1.31 (3H, t), 2.85 (2H, t), 3.65 (2H, t), 4.26 (2H, q), 4.58 (2H, s), 6.40 (1H, d, J = 16Hz), 7.15 (1H, d), 7.26 (1H, s), 7.33 (1H, d) and 7.64 (1H, d, J = 16Hz).

35 **Description 2rc**

**E-7-(2-Carboxyvinyl)-3,4-dihydro-1H-isoquinoline-2-carboxylic acid *tert* butyl ester**

A solution of 7-(2-ethoxycarbonylvinyl)-3,4-dihydro-1H-isoquinoline-2-carboxylic acid *tert* butyl ester (1.0g) in ethanol/water (30ml, 2:1) containing potassium hydroxide (0.34g) was stirred for 16h. Solvent was reduced to low volume and partitioned between ethyl acetate and water. The pH of the aqueous phase was adjusted to 1 by the addition of 5N HCl, the organic phase separated and washed with brine, dried (MgSO<sub>4</sub>) and solvent removed at reduced pressure to give 7-(2-carboxyvinyl)-3,4-dihydro-1H-isoquinoline-2-carboxylic acid *tert* butyl ester (0.76g).

MS m/z (AP<sup>+</sup>): 204 (MH<sup>+</sup>; 100%)

10

#### Description 3rc

##### E-7-Bromo-(2-Methyl-1,2,3,4-tetrahydroisoquinoline)

A solution of 7-bromo-1,2,3,4-tetrahydroisoquinoline (13.0g) in formic acid (21ml) was treated with 40% formalin and heated at 80°C for 2h. After cooling to room temperature, the reaction mixture was neutralised with sodium hydroxide (20g) and extracted with dichloromethane. The organic phase was washed with brine dried (MgSO<sub>4</sub>) and solvent removed at reduced pressure to give the title compound (13.0g) as an oil.

#### 20 Description 4rc

##### E-7-(2-Ethoxycarbonylvinyl)-2-methyl-1,2,3,4-tetrahydroisoquinoline

The title compound was prepared from 7-bromo-2-methyl-1,2,3,4-tetrahydroisoquinoline and ethyl acrylate in 30% yield according to the method of Description 1rc.

25

#### Description 5rc

##### E-7-(2-Carboxyvinyl)-2-methyl-1,2,3,4-tetrahydroisoquinoline

E-7-(2-ethoxycarbonylvinyl)-2-methyl-1,2,3,4-tetrahydroisoquinoline (4.59g) in methanol (100ml) was warmed to 50°C and sodium hydroxide (2N, 50ml) added. The mixture was stirred at 50°C for 12h, stood at room temperature for 24h and then neutralised with hydrochloric acid (2N, 50ml). Solvent was removed at reduced pressure to give a final volume of 80ml. The title compound (2.4g) crystallised on standing.

#### 35 Example 1rc

##### E-N-(4-Methoxyphenyl)-3-(1,2,3,4-tetrahydroisoquinolin-7-yl)acrylamide

(a) E-7-(2-carboxyvinyl)-3,4-dihydro-1H-isoquinoline-2-carboxylic acid *tert* butyl ester (0.47g), hydroxybenzotriazole (0.02g), 4-anisidine (0.19g) in DMF (5ml) was treated with 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide (0.30g). The



mixture was stirred for 16h, diluted with ethyl acetate, washed with water, 2N HCl, aqueous sodium carbonate and brine to give E-7-(2-carboxyvinyl)-3,4-dihydro-1H-isoquinoline-2-carboxylic acid tert butyl ester (0.59g).

m/z(AP<sup>+</sup>): 409 (MH<sup>+</sup>)

- 5 (b) A solution of E-7-[2-(4-methoxyphenylcarbamoyl)vinyl]-3,4-dihydro-1H-isoquinoline-2-carboxylic acid *tert* butyl ester (0.48g) in dichloromethane (6ml) was treated with trifluoroacetic acid and stirred for 16h. Solvent was removed at reduced pressure and the residue column chromatographed (silica gel 0 - 10% {9:1 methanol/ammonia} in dichloromethane) to give the title compound (0.24g) after
- 10 trituration with diethyl ether.

<sup>1</sup>H NMR (250MHz, CDCl<sub>3</sub>) δ; 2.82 (2H, t), 3.17 (2H, t), 3.80 (3H, s), 4.05 (2H, s), 6.67 (1H, d), 6.86 (2H, d), 7.11 (1H, d), 7.17 (1H, s), 7.33 (1H, d), 7.61 (3H, m) and 9.09 (1H, br. s); m/z(AP<sup>+</sup>): 309 (MH<sup>+</sup>; 100%)

15 **Example 2rc**

**E-N-Phenyl-3-(1,2,3,4-tetrahydroisoquinolin-7-yl)acrylamide**

- (a) From E-7-(2-carboxyvinyl)-3,4-dihydro-1H-isoquinoline-2-carboxylic acid tert butyl ester (0.48g) and aniline (0.14g), E-7-(2-phenylcarbamoylvinyl)-3,4-dihydro-1H-isoquinoline-2-carboxylic acid *tert* butyl ester (0.44g) was prepared according
- 20 to the method of Example 1rc(a).

<sup>1</sup>H NMR (250MHz, CDCl<sub>3</sub>) δ; 2.84 (2H, t), 3.65 (2H, t), 4.57 (2H, s), 6.52 (1H, d), 7.12 - 7.64 (8H, m) and 7.71 (1H, d).

- (b) From E-7-(2-phenylcarbamoylvinyl)-3,4-dihydro-1H-isoquinoline-2-carboxylic acid *tert* butyl ester (0.43g), the title compound (0.17g) isolated as a
- 25 colourless solid was prepared according to the method of Example 1rc(b).

<sup>1</sup>H NMR (250MHz, d<sup>6</sup>DMSO) δ; 2.71 (2H, t), 2.99 (2H, t), 3.90 (2H, s), 6.73 (1H, d), 7.00 (1H, t), 7.12 (1H, d), 7.24 - 7.35 (4H, m), 7.45 (1H, d), 7.63 (2H, d) and 10.12 (1H, s);

MS m/z(AP<sup>+</sup>): 279 (MH<sup>+</sup>; 100%)

30

**Example 3rc**

**E-N-Phenyl-3-(2-methyl-1,2,3,4-tetrahydroisoquinolin-7-yl)acrylamide**

- E-N-Phenyl-3-(1,2,3,4-tetrahydroisoquinolin-7-yl)acrylamide (0.13g) was heated for 150 min in a mixture of formic acid (0.54ml) and 37% formaldehyde (1ml).
- 35 After cooling the mixture was neutralised by the addition of solid sodium hydroxide and partitioned between dichloromethane and water. The organic phase was washed with 2N sodium hydroxide, water, brine and dried (MgSO<sub>4</sub>). Solvent was removed at reduced pressure and the residue column chromatographed (silica

gel 0 - 10% {9:1methanol/ammonia} in dichloromethane) to give the title compound (0.05g) as a colourless solid.

<sup>1</sup>H NMR (250MHz, CDCl<sub>3</sub>) δ; 2.47 (3H, s), 2.69 (2H, t), 2.94 (2H, t), 3.57 (2H, s), 6.50 (1H, d), 7.10 - 7.63 (8H, m) and 7.72 (1H, d); m/z(API<sup>+</sup>): 293(MH<sup>+</sup>; 100%)

5

#### Example 4rc

##### E-N-Phenyl-3-(2-benzyl-1,2,3,4-tetrahydroisoquinolin-7-yl)acrylamide

E-N-phenyl-3-(1,2,3,4-tetrahydroisoquinolin-7-yl)acrylamide (0.28g), benzaldehyde (0.1ml), acetic acid (0.057ml) and methanol (20ml) were combined and treated with sodium cyanoborohydride (0.63g). The mixture was stirred for 16h, solvent was removed *in vacuo* and the residue column was chromatographed (silica gel 0 - 10% {9:1 methanol-ammonia} in dichloromethane) to give the title compound (0.23g) as a colourless foam.

<sup>1</sup>H NMR (250MHz, CDCl<sub>3</sub>) δ; 2.79 (2H, t), 2.95 (2H, t), 3.65 (2H, s), 3.73 (2H, s), 6.48 (1H, d), 7.10 - 7.63 (11H, m) 7.62 (2H, d,) and 7.69 (1H, d); m/z(API<sup>+</sup>): 369(MH<sup>+</sup>; 100%)

#### Example 5rc

##### E-N-Phenyl-3-(2-n-propyl-1,2,3,4-tetrahydroisoquinolin-7-yl)acrylamide

From E-N-phenyl-3-(1,2,3,4-tetrahydroisoquinolin-7-yl)acrylamide (0.28g) and propionaldehyde (0.07ml), the title compound (0.22g) isolated as a colourless solid was prepared according to the method of Example 4rc.

<sup>1</sup>H NMR (250MHz, CDCl<sub>3</sub>) δ; 1.05 (3H, t), 1.78 (2H, m), 2.87 (2H, t), 3.06 (2H, d), 3.14 (2H, d), 4.13 (2H, s), 6.80 (1H, d), 7.05 - 7.37 (6H, m), 7.58 (1H, d), 7.75 (2H, d) and 9.24 (1H, s); m/z(API<sup>+</sup>): 321(MH<sup>+</sup>; 100%)

#### Example 6rc

##### E-N-Phenyl-3-(2-ethyl-1,2,3,4-tetrahydroisoquinolin-7-yl)acrylamide

From E-N-phenyl-3-(1,2,3,4-tetrahydroisoquinolin-7-yl)acrylamide (0.28g) and acetaldehyde (0.06ml), the title compound (0.88g) isolated as a colourless solid was prepared according to the method of Example 4rc.

<sup>1</sup>H NMR (250MHz, CDCl<sub>3</sub>) δ; 1.20 (3H, t), 2.60 (2H, q), 2.74 (2H, t), 2.93 (2H, t), 3.62 (2H, s), 6.50 (1H, d), 7.12 (2H, t), 7.19 (1H, s), 7.26 - 7.38 (3H, m) 7.61 (2H, d) and 7.69(1H, d); m/z(API<sup>+</sup>): 307(MH<sup>+</sup>; 100%)

35

#### Example 7rc

##### E-N-(3-Cyanophenyl)-3-(1,2,3,4-tetrahydroisoquinolin-7-yl)acrylamide

(a) From E-7-(2-carboxyvinyl)-3,4-dihydro-1H-isoquinoline-2-carboxylic acid tert butyl ester (1.01g) and 3-aminobenzonitrile (0.39g), E-7-[2-(3-

cyanophenyl)carbamoylvinyl]-3,4-dihydro-1H-isoquinoline-2-carboxylic acid *tert* butyl ester (0.41g) was prepared according to the method of Example 1rc(a).

$m/z$ (API<sup>+</sup>): 304 (MH<sup>+</sup> - tertbutoxycarbonyl; 100%)

- 5 (b) From E-7-[2-(3-cyanophenyl)carbamoylvinyl]-3,4-dihydro-1H-isoquinoline-2-carboxylic acid *tert* butyl ester (0.38g), the title compound (0.08g) isolated as a colourless solid was prepared according to the method of Example 1rc(b).

<sup>1</sup>H NMR (250MHz, d<sup>6</sup>DMSO)  $\delta$ : 2.73(t, 2H), 2.98(t, 2H), 3.90(s, 2H), 6.75(d, 1H), 7.15(d, 1H), 7.30(s, 1H), 7.39(d, 1H), 7.58(d, 1H), 7.85(m, 1H), 8.25(s, 1H) and 10.51(s, 1H);

- 10 MS  $m/z$ (API<sup>+</sup>): 304 (MH<sup>+</sup>; 100%)

#### Example 8rc

**E-N-(3-Cyanophenyl)-3-(2-methyl-1,2,3,4-tetrahydroisoquinolin-7-yl)acrylamide**

- 15 From E-N-(3-cyanophenyl)-3-(1,2,3,4-tetrahydroisoquinolin-7-yl)acrylamide (0.06g) the title compound (0.03g), isolated as a colourless solid was prepared according to the method of Example 3rc.

<sup>1</sup>H NMR (250MHz, d<sup>6</sup>DMSO)  $\delta$ : 2.35 (3H, s), 2.60 (2H, t), 2.84 (2H, t), 3.51 (2H, s), 6.74 (1H, d), 7.18 (1H, d), 7.31 (1H, s), 7.40 (1H, d), 7.51 - 7.59 (3H, m) 7.87 (1H, d), 8.21 (1H, s) and 10.49 (1H, s). (1H, d), 7.85 (1H, m), 8.25 (1H, s) and 10.51 (1H, s);

MS  $m/z$ (API<sup>+</sup>): 318 (MH<sup>+</sup>; 100%)

#### Example 9rc

- 25 **E-N-(2-Chlorophenyl)-3-(2-methyl-1,2,3,4-tetrahydroisoquinolin-7-yl)acrylamide**

From E-E-7-(2-carboxyvinyl)-2-methyl-1,2,3,4-tetrahydroisoquinoline (0.22g) and 2-chloroaniline (0.13g) the title compound (0.03g) was prepared according to the method of Example 1rc(a) omitting the acid wash.

- 30 <sup>1</sup>H NMR (250MHz, CDCl<sub>3</sub>)  $\delta$ : 2.50 (3H, s), 2.74 (2H, t), 2.96 (2H, t), 3.63 (2H, s), 6.54 (1H, d), 7.06 (1H, t), 7.15 (1H, d), 7.23 - 7.46 (4H, m), 7.71 (1H, d), 7.78 (1H, br. s.) and 8.54 (1H, d);  $m/z$ (API<sup>+</sup>): 327(MH<sup>+</sup>; 100%)

#### Example 10rc

- 35 **E-N-(2-Methoxyphenyl)-3-(2-methyl-1,2,3,4-tetrahydroisoquinolin-7-yl)acrylamide**

From E-7-(2-carboxyvinyl)-2-methyl-1,2,3,4-tetrahydroisoquinoline (0.22g) and 2-methoxyaniline (0.12g), the title compound (0.13g) was prepared according to the method of Example 1rc(a) omitting the acid wash.

<sup>1</sup>H NMR (250MHz, d<sup>6</sup>DMSO) δ; 2.44 (3H, s), 2.66 (2H, t), 2.91 (2H, t), 3.55 (2H, s), 3.87 (3H, s), 6.55 (1H, d), 6.86 (1H, dd), 6.93 - 7.07 (3H, m), 7.17 (1H, s), 7.29 (1H, t), 7.66 (1H, d), 7.99 (1H, br. s.) and 8.50 (1H, d); m/z(API<sup>+</sup>): 323 (MH<sup>+</sup>; 100%)

5

**Example 11rc****E-N-(3-Methoxyphenyl)-3-(2-methyl-1,2,3,4-tetrahydroisoquinolin-7-yl)acrylamide**

From E-7-(2-carboxyvinyl)-2-methyl-1,2,3,4-tetrahydroisoquinoline (0.22g) and 3-methoxyaniline (0.12g), the title compound (0.10g) was prepared according to the method of Example 1rc(a) omitting the acid wash.

<sup>1</sup>H NMR (250MHz, CDCl<sub>3</sub>) δ; 2.44 (3H, s), 2.67 (2H, m), 2.90 (2H, m), 3.51 (2H, s), 3.78 (3H, s), 6.54 (1H, d), 6.67 (1H, d), 7.02 - 7.26 (5H, m), 7.46 (1H, s), 7.66 (1H, d) and 7.98 (1H, s); m/z(API<sup>+</sup>): 322 (MH<sup>+</sup>; 100%)

15

**Example 12rc****E-N-(3-Chlorophenyl)-3-(2-methyl-1,2,3,4-tetrahydroisoquinolin-7-yl)acrylamide**

From E-7-(2-carboxyvinyl)-2-methyl-1,2,3,4-tetrahydroisoquinoline (0.22g) and 3-chloroaniline (0.13g), the title compound (0.09g) was prepared according to the method of Example 1rc(a) omitting the acid wash.

<sup>1</sup>H NMR (250MHz, CDCl<sub>3</sub>) δ; 2.79 (3H, s), 3.08 (2H, m), 3.16 (2H, m), 4.06 (2H, s), 6.47 (1H, d), 6.95 (1H, d), 6.97 (1H, s), 7.14 (2H, dt), 7.20 - 7.29 (1H, m), 7.37 (1H, d), 7.69 (1H, m), 7.94 (1H, m) and 9.43 (1H, br. s.); m/z(API<sup>+</sup>): 327 (MH<sup>+</sup>; 100%)

25

**Example 13rc****E-N-(4-Chlorophenyl)-3-(2-methyl-1,2,3,4-tetrahydroisoquinolin-7-yl)acrylamide**

From E-7-(2-carboxyvinyl)-2-methyl-1,2,3,4-tetrahydroisoquinoline (0.22g) and 4-chloroaniline, the title compound (0.09g) was prepared according to the method of Example 1rc(a) omitting the acid wash.

<sup>1</sup>H NMR (250MHz, d<sup>6</sup>DMSO) δ; 2.41 (3H, s), 2.69 (2H, m), 2.86 (2H, m), 3.59 (2H, s), 6.76 (1H, d), 7.18 (1H, d), 7.30 (1H, s), 7.37 - 7.40 (3H, m), 7.54 (1H, d), 7.79 (2H, d) and 10.34 (1H, br. s); m/z(API<sup>+</sup>): 327 (MH<sup>+</sup>; 100%)

35

**Example 14rc****E-N-Methyl-N-benzyl-3-(2-methyl-1,2,3,4-tetrahydroisoquinolin-7-yl)acrylamide**

From E-7-(2-carboxyvinyl)-2-methyl-1,2,3,4-tetrahydroisoquinoline (0.22g) and N-methylbenzylamine (0.12g), the title compound (0.16g) was prepared according to the method of Example 1rc(a) omitting the acid wash.

- 5 <sup>1</sup>H NMR (250MHz, CDCl<sub>3</sub>) δ; 2.46 (3H, d), 2.70 (2H, m), 2.91 (2H, m), 3.07 (3H, d), 3.57 (2H, d), 4.69 (2H, d), 6.85 (1H, t), 7.06 - 7.38 (8H, m) and 7.72 (1H, d); MS m/z(API<sup>+</sup>): 320 (MH<sup>+</sup>; 100%)

#### Example 15rc

- 10 **E-N-(3-Nitrophenyl)-3-(2-methyl-1,2,3,4-tetrahydroisoquinolin-7-yl)acrylamide**

From E-7-(2-carboxyvinyl)-2-methyl-1,2,3,4-tetrahydroisoquinoline (0.22g) and 3-nitroaniline (0.14g), the title compound (0.04g) was prepared according to the method of Example 1rc(a) omitting the acid wash.

- 15 <sup>1</sup>H NMR (250MHz, CDCl<sub>3</sub>) δ; 2.46(3H, s), 2.73 (2H, m), 2.89 (2H, m), 3.54 (2H, s), 4.69 (2H, d), 6.54 (1H, d), 7.03 (2H, m), 7.19 (1H, d), 7.43 (1H, t), 7.64 (1H, d), 7.92 (1H, d), 8.10 (1H, d), 8.48 (1H, s) and 8.88 (1H, br s.); m/z(API<sup>+</sup>): 338 (MH<sup>+</sup>; 100%)

#### Example 16rc

- 20 **E-N-Methyl-N-phenyl-3-(2-methyl-1,2,3,4-tetrahydroisoquinolin-7-yl)acrylamide**

From E-7-(2-carboxyvinyl)-2-methyl-1,2,3,4-tetrahydroisoquinoline (0.22g) and N-methylaniline (0.11g), the title compound (0.15g) was prepared according to the method of Example 1rc(a) omitting the acid wash.

- 25 <sup>1</sup>H NMR (250MHz, CDCl<sub>3</sub>) δ; 2.42 (3H, s), 2.64 (2H, t), 2.87 (2H, t), 3.40 (3H, s), 3.50 (2H, s), 6.30 (1H, d), 6.96 - 7.10 (2H, m), 7.21 - 7.26 (2H, m), 7.34 - 7.49 (2H, m) and 7.62 (1H, d); m/z(API<sup>+</sup>): 307 (MH<sup>+</sup>; 100%)

#### Example 17rc

- 30 **E-N-(3-Carbomethoxyphenyl)-3-(2-methyl-1,2,3,4-tetrahydroisoquinolin-7-yl)acrylamide**

From E-7-(2-carboxyvinyl)-2-methyl-1,2,3,4-tetrahydroisoquinoline (0.22g) and 3-carbomethoxyaniline (0.15g), the title compound (0.07g) was prepared according to the method of Example 1rc(a) omitting the acid wash.

- 35 <sup>1</sup>H NMR (250MHz, CDCl<sub>3</sub>) δ; 2.57 (3H, s), 2.79 (2H, m), 2.96 (2H, m), 3.73 (2H, s), 3.91 (3H, s), 6.49 (1H, d), 7.04 (1H, d), 7.08 (1H, s), 7.20 (1H, d), 7.45 (1H, t), 7.59 (1H, d), 7.78 (1H, d), 8.06 (1H, d), 8.245 (1H, s) and 8.50 (1H, s); m/z(API<sup>+</sup>): 351 (MH<sup>+</sup>; 100%)

**Example 18rc****E-N-Methyl-3-[3-(2-methyl-1,2,3,4-tetrahydroisoquinolin-7-yl)acryloylamino]benzamide**

From E-7-(2-carboxyvinyl)-2-methyl-1,2,3,4-tetrahydroisoquinoline (0.22g) and  
5 3-N-methylcarboxamidoaniline (0.15g), the title compound (0.07g) was prepared according to the method of Example 1rc(a) omitting the acid wash.

<sup>1</sup>H NMR (250MHz, d<sup>6</sup>-DMSO) δ; 2.79 (3H, d), 2.83 - 2.94 (4H, m), 3.17 (3H, s), 3.72 (2H, s), 6.56 and 6.84 (1H, d), 6.99 (1H, t), 7.14 - 7.59 (5H, m), 7.88 (1H, t), 8.13 (1H, s), 8.44 (1H, m) and 10.46 (1H, s); m/z(API<sup>+</sup>): 350 (MH<sup>+</sup>)

10

**Example 19rc****E-N-(3-N-Methylsulphonylphenyl)-3-(2-methyl-1,2,3,4-tetrahydroisoquinolin-7-yl) acrylamide**

A solution of E-7-(2-carboxyvinyl)-2-methyl-1,2,3,4-tetrahydroisoquinoline  
15 (0.22g) in dichloromethane (10ml) was treated with oxalyl chloride (0.30ml) and dimethylformamide (2 drops). The mixture was stirred for 2h, solvent removed at reduced pressure and the residue treated sequentially with 3-methylsulphonylaniline hydrochloride (0.21g), tetrahydrofuran (20ml) and triethylamine (1ml). The reaction mixture was stirred for 16h, diluted with ethyl  
20 acetate and washed with water. The organic phase was dried (MgSO<sub>4</sub>) and solvent removed at reduced pressure. The residue was column chromatographed (silica gel, 5% methanol/dichloromethane) to give the title compound (0.07g).

<sup>1</sup>H NMR (250MHz, d<sup>6</sup>-DMSO with D<sub>2</sub>O shake) δ; 2.69 (3H, s), 3.03 (2H, m), 3.12 (2H, m), 3.20 (3H, s), 4.03 (2H, s), 6.81 (1H, d), 7.30 (1H, d), 7.43 (1H, s),  
25 7.53 - 7.65 (4H, m), 7.91 (1H, br. s) and 8.38 (1H, s); m/z(API<sup>+</sup>): 371 (MH<sup>+</sup>)

**Example 20rc****E-N-(1,3-Oxazol-5-ylphenyl)-3-(2-methyl-1,2,3,4-tetrahydroisoquinolin-7-yl)acrylamide**

30 From E-7-(2-carboxyvinyl)-2-methyl-1,2,3,4-tetrahydroisoquinoline (0.22g) and 1,3-oxazol-5-ylaniline (0.16g), the title compound (0.15g) was prepared according to the method of Example 19rc omitting triethylamine.

<sup>1</sup>H NMR (250MHz, CDCl<sub>3</sub>) δ; 2.43 (3H, s), 2.67 (2H, t), 2.91 (2H, t), 3.51 (2H, s), 6.57 (1H, d), 7.04 - 7.10 (2H, m), 7.24 - 7.37 (3H, m), 7.60 (1H, m), 7.73 (1H, d),  
35 7.84 (1H, s), 8.03 (1H, s) and 8.17 (1H, s); m/z(API<sup>+</sup>): 360 (MH<sup>+</sup>; 100%)

**Example 21rc****E-N-(3-Acetylaminophenyl)-3-(2-methyl-1,2,3,4-tetrahydroisoquinolin-7-yl)acrylamide**

From E-7-(2-carboxyvinyl)-2-methyl-1,2,3,4-tetrahydroisoquinoline (0.22g) and 3-acetylaminoaniline (0.15g), the title compound (0.13g) was prepared according to the method of Example 19rc omitting triethylamine.

- 5 <sup>1</sup>H NMR (250MHz, CDCl<sub>3</sub>). 2.04 (3H, s), 2.39 (3H, s), 2.63 (2H, m), 2.86 (2H, m), 3.49 (2H, s), 6.54 (1H, d), 6.96 - 7.37 (5H, m), 7.56 (1H, d), 7.82 (1H, s), 8.27 (1H, s) and 8.65 (1H, s); m/z(API<sup>+</sup>): 350 (MH<sup>+</sup>; 100%).

#### Example 22rc

- 10 **E-N-(3-Ethylphenyl)-3-(2-methyl-1,2,3,4-tetrahydroisoquinolin-7-yl)acrylamide**

From E-7-(2-carboxyvinyl)-2-methyl-1,2,3,4-tetrahydroisoquinoline (0.22g) and 3-ethylaniline (0.12g), the title compound (0.10g) was prepared according to the method of Example 19rc omitting triethylamine.

- 15 <sup>1</sup>H NMR (250MHz, CDCl<sub>3</sub>) δ: 1.20 (3H, t), 2.43 (3H, s), 2.60 (2H, q), 2.67 (2H, m), 2.90 (2H, m), 3.51 (2H, s), 6.55 (1H, d), 6.94 (1H, d), 7.04 - 7.26 (4H, m), 7.42 - 7.50 (2H, m), 7.68 (1H, d) and 7.89 (1H, s); m/z(API<sup>+</sup>): 321 (MH<sup>+</sup>; 100%)

#### Example 23rc

- 20 **E-N-(3-Methylphenyl)-3-(2-methyl-1,2,3,4-tetrahydroisoquinolin-7-yl)acrylamide**

From E-7-(2-carboxyvinyl)-2-methyl-1,2,3,4-tetrahydroisoquinoline (0.22g) and 3-methylaniline (0.12g), the title compound (0.09g) was prepared according to the method of Example 19rc omitting triethylamine.

- 25 <sup>1</sup>H NMR (250MHz, CDCl<sub>3</sub>) δ 2.32 (3H, s), 2.45 (3H, s), 2.68 (2H, t), 2.91 (2H, m), 3.52 (2H, s), 6.53 (1H, d), 6.91 (1H, d), 7.05 - 7.28 (4H, m), 7.38 (1H, d), 7.48 (1H, s), 7.68 (1H, d) and 7.69 (1H, s); m/z(API<sup>+</sup>): 307 (MH<sup>+</sup>; 100%)

#### Example 24rc

- 30 **E-N-(3-*tert*-Butylphenyl)-3-(2-methyl-1,2,3,4-tetrahydroisoquinolin-7-yl)acrylamide**

From E-7-(2-carboxyvinyl)-2-methyl-1,2,3,4-tetrahydroisoquinoline (0.22g) and 3-*tert*-butylaniline (0.15g), the title compound (0.11g) was prepared according to the method of Example 19rc omitting triethylamine.

- 35 <sup>1</sup>H NMR (250MHz, CDCl<sub>3</sub>) δ: 1.28 (9H, s), 2.42 (3H, s), 2.65 (2H, t), 2.89 (2H, m), 3.48 (2H, s), 6.60 (1H, d), 6.91 (1H, d), 7.01 - 7.23 (4H, m), 7.64 - 7.69 (2H, m), 8.22 (1H, s);

MS m/z(API<sup>+</sup>): 349 (MH<sup>+</sup>; 100%)

#### Example 25rc

**E-N-(4-Fluorophenyl)-3-(2-methyl-1,2,3,4-tetrahydroisoquinolin-7-yl)acrylamide**

From E-7-(2-carboxyvinyl)-2-methyl-1,2,3,4-tetrahydroisoquinoline (0.22g) and 4-fluoroaniline (0.12g), the title compound (0.15g) was prepared according to the method of Example 19rc omitting triethylamine.

<sup>1</sup>H NMR (250MHz, CDCl<sub>3</sub>) δ; 2.44 (3H, s), 2.67 (2H, t), 2.91 (2H, t), 3.52 (2H, s), 6.50 (1H, d), 6.80 - 7.10 (4H, m), 7.25 (1H, d), 7.41 (2H, br.m.), 7.67 (1H, d) and 7.82 (1H, br. s); m/z(API<sup>+</sup>): 311 (MH<sup>+</sup>; 100%)

**Example 26rc****E-N-(4-Methoxyphenyl)-3-(2-methyl-1,2,3,4-tetrahydroisoquinolin-7-yl)acrylamide**

From E-7-(2-carboxyvinyl)-2-methyl-1,2,3,4-tetrahydroisoquinoline (0.22g) and 4-methoxyaniline (0.12g), the title compound (0.15g) was prepared according to the method of Example 19rc omitting triethylamine.

<sup>1</sup>H NMR (250MHz, CDCl<sub>3</sub>) δ; 2.45 (3H, s), 2.68 (2H, t), 2.92 (2H, t), 3.55 (2H, s), 3.79 (3H, s), 6.49 (1H, d), 6.87 (2H, d), 7.11 - 7.63 (3H, d), 7.52 (2H, br.s) and 7.66 (1H, d);

MS m/z(API<sup>+</sup>): 323 (MH<sup>+</sup>; 100%)

**Example 27rc****E-N-(4-Carbomethoxyphenyl)-3-(2-methyl-1,2,3,4-tetrahydroisoquinolin-7-yl)acrylamide**

From E-7-(2-carboxyvinyl)-2-methyl-1,2,3,4-tetrahydroisoquinoline (0.22g) and 4-carbomethoxyaniline (0.15g), the title compound (0.19g) was prepared according to the method of Example 19rc omitting triethylamine.

<sup>1</sup>H NMR (250MHz, CDCl<sub>3</sub>) δ; 2.46 (3H, s), 2.69 (2H, t), 2.93 (2H, t), 3.55 (2H, s), 3.90 (3H, s), 6.52 (1H, d), 7.09 - 7.31 (4H, m), 7.68 - 7.73 (2H, m), 7.80 (1H, d) and 8.01 (2H, d); m/z(API<sup>+</sup>): 351 (MH<sup>+</sup>; 100%)

**Example 28rc****E-N-(4-Cyanophenyl)-3-(2-methyl-1,2,3,4-tetrahydroisoquinolin-7-yl)acrylamide**

From E-7-(2-carboxyvinyl)-2-methyl-1,2,3,4-tetrahydroisoquinoline (0.22g) and 4-aminobenzonitrile (0.12g), the title compound (0.03g) was prepared according to the method of Example 19rc omitting triethylamine.

<sup>1</sup>H NMR (250MHz, CDCl<sub>3</sub>) δ; 2.50 (3H, s), 2.75 (2H, t), 2.96 (2H, t), 3.61 (2H, s), 6.52 (1H, d), 7.15 (1H, d), 7.26 (1H, s), 7.30 (1H, d), 7.61 (2H, d), 7.69 (1H, d), 7.78 (2H, d) and 7.95 (1H, s); m/z(API<sup>+</sup>): 318 (MH<sup>+</sup>; 100%)



**Example 29rc****E-N-(4-Nitrophenyl)-3-(2-methyl-1,2,3,4-tetrahydroisoquinolin-7-yl)acrylamide**

- 5 From E-7-(2-carboxyvinyl)-2-methyl-1,2,3,4-tetrahydroisoquinoline (0.22g) and 4-nitroaniline (0.14g), the title compound (0.09g) was prepared according to the method of Example 19rc omitting triethylamine.

10 <sup>1</sup>H NMR (250MHz, d<sup>6</sup>-DMSO) δ; 2.50 (3H, s), 2.60 (2H, t), 2.82 (2H, m), 3.50 (2H, s), 6.47 (1H, d), 7.18 (1H, d), 7.32 (1H, s), 7.40 (1H, d), 7.64 (1H, d), 7.93 (2H, d), 8.24 (2H, d), 10.76 (1H, br. s); m/z(API<sup>+</sup>): 337 (MH<sup>+</sup>; 100%)

**Example 30rc****E-N-(4-Methylphenyl)-3-(2-methyl-1,2,3,4-tetrahydroisoquinolin-7-yl)acrylamide**

- 15 From E-7-(2-carboxyvinyl)-2-methyl-1,2,3,4-tetrahydroisoquinoline (0.22g) and 4-toluidine (0.11g), the title compound (0.09g) was prepared according to the method of Example 19rc omitting triethylamine.

20 <sup>1</sup>H NMR (250MHz, d<sup>6</sup>-DMSO) 2.31 (3H, s), 2.44 (3H, s), 2.67 (2H, m), 2.91 (2H, m), 3.51 (2H, s), 6.53 (1H, d), 6.99 - 7.13 (4H, m), 7.24 (1H, d), 7.51 (2H, d), 7.66 (1H, d) and 7.92 (1H, d); m/z(API<sup>+</sup>): 306 (MH<sup>+</sup>; 100%)

**Example 31rc****E-N-(3-Methoxy-5-trifluoromethylphenyl)-3-(2-methyl-1,2,3,4-tetrahydroisoquinolin-7-yl)acrylamide**

- 25 From E-7-(2-carboxyvinyl)-2-methyl-1,2,3,4-tetrahydroisoquinoline (0.22g) and 3-methoxy-5-trifluoromethylaniline (0.19g), the title compound (0.16g) was prepared according to the method of Example 19rc.

30 <sup>1</sup>H NMR (250MHz, CDCl<sub>3</sub>) δ; 2.46 (3H, s), 2.67 (2H, t), 2.93 (2H, m), 3.56 (2H, s), 3.83 (3H, s), 6.49 (1H, d), 6.88 (1H, s), 7.10 (1H, d), 7.15 (1H, s), 7.26 - 7.31 (2H, m), 7.66 (1H, d) and 7.73 (1H, d); m/z(API<sup>+</sup>): 390 (MH<sup>+</sup>; 100%)

**Example 32rc****E-1-(3,4-Dihydro-1H-isoquinolin-2-yl)-3-(2-methyl-1,2,3,4-tetrahydroisoquinolin-7-yl)propenone**

- 35 From E-7-(2-carboxyvinyl)-2-methyl-1,2,3,4-tetrahydroisoquinoline (0.22g) and 1,2,3,4-tetrahydroisoquinoline (0.13g) the title compound (0.07g) was prepared according to the method of Example 1rc(a) omitting the acid wash.

<sup>1</sup>H NMR (250MHz, CDCl<sub>3</sub>) δ; 2.47 (3H, s), 2.70 (2H, t), 2.94 (4H, m), 3.60 (2H, s), 3.88 (2H, m), 6.89 (1H, d), 7.11 - 7.21 (6H, m), 7.34 (1H, d) and 7.67 (1H, d);

m/z(API<sup>+</sup>): 333(MH<sup>+</sup>; 100%)

#### Example 33rc

##### **E-1-(3,4-Dihydro-2H-quinolin-1-yl)-3-(2-methyl-1,2,3,4-tetrahydroisoquinolin-7-yl)propenone**

From E-7-(2-carboxyvinyl)-2-methyl-1,2,3,4-tetrahydroisoquinoline (0.22g) and 1,2,3,4-tetrahydroquinoline (0.13g) the title compound (0.07g) was prepared according to the method of Example 1rc(a) omitting the acid wash.

<sup>1</sup>H NMR (250MHz, d<sup>6</sup>-DMSO) 1.69 (2H, t), 2.11 (3H, s), 2.33 (2H, m), 2.37 (2H, m), 2.52 (2H, t), 3.23 (2H, s), 3.59 (2H, t), 6.64 (1H, d), 6.91 - 7.10 (7H, m) and 7.31 (1H, d);

MS m/z(API<sup>+</sup>): 333(MH<sup>+</sup>; 100%)

#### Example 34rc

##### **E-1-(3,3-Dimethyl-2,3-dihydroindol-1-yl)-3-(2-methyl-1,2,3,4-tetrahydroisoquinolin-7-yl)propenone**

From E-7-(2-carboxyvinyl)-2-methyl-1,2,3,4-tetrahydroisoquinoline (0.22g) and 3,3-dimethylindoline (0.15g) the title compound (0.06g) was prepared according to the method of Example 1rc(a) omitting the acid wash.

<sup>1</sup>H NMR (250MHz, CDCl<sub>3</sub>) 1.39 (6H, s), 2.47 (3H, s), 2.70 (2H, t), 2.92 (2H, m), 3.61 (2H, s), 4.00 (2H, s), 6.78 (1H, d), 7.04 - 7.26 (5H, m), 7.36 (1H, d), 7.78 (1H, d) and 8.31 (1H, br. s.); m/z(API<sup>+</sup>): 346(MH<sup>+</sup>; 100%)

#### Example 35rc

##### **E-1-(2,3-Dihydroindol-1-yl)-3-(2-methyl-1,2,3,4-tetrahydroisoquinolin-7-yl)propenone**

From E-7-(2-carboxyvinyl)-2-methyl-1,2,3,4-tetrahydroisoquinoline (0.22g) and indoline (0.12g) the title compound (0.11g) was prepared according to the method of Example 1rc(a) omitting the acid wash.

<sup>1</sup>H NMR (250MHz, CDCl<sub>3</sub>) 2.48 (3H, s), 2.71 (2H, t), 2.95 (2H, t), 3.61 (2H, s), 4.28 (2H, t), 6.84 (1H, d), 7.03 (1H, t), 7.13 (1H, d), 7.20 (3H, m), 7.35 (1H, d), 7.78 (1H, d) and 8.36 (1H, br s); m/z(API<sup>+</sup>): 318(MH<sup>+</sup>; 100%)

### PHARMACOLOGICAL DATA

#### **1. Binding Assay Method**

International Application Publication Number WO 92/22293 (SmithKline Beecham) discloses compounds having anti-convulsant activity, including *inter alia* the compound *trans*-(+)-6-acetyl-4S-(4-fluorobenzoylamino)-3,4-dihydro-2,2-dimethyl-2H-1-benzopyran-3R-ol (hereinafter referred to as Compound A). It has

been found that the compounds of WO 92/22293 bind to a novel receptor obtainable from rat forebrain tissue, as described in WO 96/18650 (SmithKline Beecham). The affinity of test compounds to the novel receptor site is assessed as follows.

5

### Method

Whole forebrain tissue is obtained from rats. The tissue is first homogenised in buffer (usually 50mM Tris/HCl, pH 7.4). The homogenised tissue is washed by centrifugation and resuspension in the same buffer, then stored at -70°C until used.

10 To carry out the radioligand binding assay, aliquots of tissue prepared as above (usually at a concentration of 1-2mg protein/ml) are mixed with aliquots of [3H]-Compound A dissolved in buffer. The final concentration of [3H]-Compound A in the mixture is usually 20nM. The mixture is incubated at room temperature for 1 hour. [3H]-Compound A bound to the tissue is then separated  
15 from unbound [3H]-Compound A by filtration through Whatman GF/B glass fibre filters. The filters are then washed rapidly with ice-cold buffer. The amount of radioactivity bound to the tissue trapped on the filters is measured by addition of liquid scintillation cocktail to the filters followed by counting in a liquid scintillation counter.

20 In order to determine the amount of "specific" binding of [3H]-Compound A, parallel assays are carried out as above in which [3H]-Compound A and tissue are incubated together in the presence of unlabelled Compound A (usually 3 µM). The amount of binding of [3H]-Compound A remaining in the presence of this unlabelled compound is defined as "non-specific" binding. This amount is  
25 subtracted from the total amount of [3H]-Compound A binding (i.e. that present in the absence of unlabelled compound) to obtain the amount of "specific" binding of [3H]-Compound A to the novel site.

The affinity of the binding of test compounds to the novel site can be estimated by incubating together [3H]-Compound A and tissue in the presence of  
30 a range of concentrations of the compound to be tested. The decrease in the level of specific [3H]-Compound A binding as a result of competition by increasing concentrations of the compound under test is plotted graphically, and non-linear regression analysis of the resultant curve is used to provide an estimate of compound affinity in terms of pKi value.

35

### Results

Compounds of this invention were active in this test with pKi values greater than 6. For example, compounds of Examples 9c, 27c and 2rc gave pKi values greater than 7.5.

## 2. MEST Test

The maximal electroshock seizure (MEST) threshold test in rodents is particularly sensitive for detecting potential anticonvulsant properties<sup>1</sup>. In this model, anticonvulsant agents elevate the threshold to electrically-induced seizures whilst proconvulsants lower the seizure threshold.

### Method

Mice (naive male, Charles River, U.K. CD-1 strain, 25 - 30g) are randomly assigned to groups of 10 - 20 and dosed orally or intraperitoneally at a dose volume of 10 ml/kg with various doses of compound (0.3 - 300 mg/kg) or vehicle. Mice are then subjected at 30 or 60 min post dose to a single electroshock (0.1 sec, 50Hz, sine wave form) administered via corneal electrodes. The mean current and standard error required to induce a tonic seizure in 50% (CC<sub>50</sub>) of the mice in a particular treatment group is determined by the 'up and down' method of Dixon and Mood (1948)<sup>2</sup>. Statistical comparisons between vehicle- and drug-treated groups are made using the method of Litchfield and Wilcoxon (1949)<sup>3</sup>.

In control animals the CC<sub>50</sub> is usually 14 - 18 mA. Hence the first animal in the control group is subjected to a current of 16 mA. If a tonic seizure does not ensue, the current is increased for a subsequent mouse. If a tonic convulsion does occur, then the current is decreased, and so on until all the animals in the group have been tested.

The percentage increase or decrease in CC<sub>50</sub> for each group compared to the control is calculated.

Studies are carried out using a Hugo Sachs Elektronik Constant Current Shock Generator with totally variable control of shock level from 0 to 300 mA and steps of 2 mA are usually used.

Drugs are suspended in 1% methyl cellulose.

### References

1. Loscher, W. and Schmidt, D. (1988). *Epilepsy Res.*, **2**, 145-181
2. Dixon, W.J. and Mood, A.M. (1948). *J. Amer. Stat. Assn.*, **43**, 109-126
3. Litchfield, J.T. and Wilcoxon, F. (1949). *J. Pharmacol. exp. Ther.*, **96**, 99-113


### Results

Compounds of this invention dosed by the oral route as a suspension in methyl cellulose and tested one hour post dosing show an increase in seizure threshold. For example, at a dose of 10 mg/kg p.o. the compounds of Examples 9c, 27c and

*2rc* showed statistically significant increases of 245, 192 and 140 % respectively.

# Claims

1. Accordingly, the present invention provides a compound of formula (I) or pharmaceutically acceptable salt thereof:



in which

- 10 Z is a carbocyclic or heterocyclic or a fused carbocyclic or heterocyclic ring containing at least one aromatic ring;  
 X is CH or N;  
 Y is hydrogen, C<sub>1-6</sub>alkyl, or a halogen;  
 P is -CH=CH- and Q is -NR<sup>1</sup>-, or;  
 P is -CH=CH- and Q is -NR<sup>1</sup>CH<sub>2</sub>-, or;  
 15 P is -NH- and Q is -CR<sup>1a</sup>=CH-;  
 R<sup>1</sup> is hydrogen, phenylC<sub>1-6</sub> alkyl, or C<sub>1-6</sub> alkyl;  
 R<sup>1a</sup> is hydrogen, halogen, phenylC<sub>1-6</sub> alkyl, or C<sub>1-6</sub> alkyl;  
 R<sup>2</sup> is hydrogen or up to three substituents selected from halogen, NO<sub>2</sub>, CN, N<sub>3</sub>, CF<sub>3</sub>O-, CF<sub>3</sub>S-, CF<sub>3</sub>CO-, CF<sub>3</sub>SO<sub>2</sub>, trifluoromethyldiaziriny, C<sub>1-6</sub>alkyl,  
 20 C<sub>1-6</sub>alkenyl, C<sub>1-6</sub>alkynyl, C<sub>1-6</sub>perfluoroalkyl, C<sub>3-6</sub>cycloalkyl, C<sub>3-6</sub>cycloalkyl-C<sub>1-4</sub>alkyl-, C<sub>1-6</sub>alkylO-, C<sub>1-6</sub>alkylCO-, C<sub>3-6</sub>cycloalkylO-, C<sub>3-6</sub>cycloalkylCO-, C<sub>3-6</sub>cycloalkyl-C<sub>1-4</sub>alkylO-, C<sub>3-6</sub>cycloalkyl-C<sub>1-4</sub>alkylCO-, phenyl, phenoxy, benzyloxy, benzoyl, phenyl-C<sub>1-4</sub>alkyl-, C<sub>1-6</sub>alkylS-, C<sub>1-6</sub>alkylSO<sub>2</sub>-, or 1,3-oxazol-5-yl(C<sub>1-4</sub>alkyl)<sub>2</sub>NSO<sub>2</sub>-, (C<sub>1-4</sub>alkyl)NHSO<sub>2</sub>-,  
 25 (C<sub>1-4</sub>alkyl)<sub>2</sub>NCO-, (C<sub>1-4</sub>alkyl)NHCO- or CONR<sup>4</sup>R<sup>5</sup>, CO<sub>2</sub>R<sup>4</sup>, or -NR<sup>4</sup>R<sup>6</sup> or NHCOR<sup>4</sup>  
 where R<sup>4</sup> and R<sup>5</sup> are each independently hydrogen or C<sub>1-4</sub> alkyl, and;  
 R<sup>6</sup> is hydrogen, C<sub>1-4</sub>alkyl, formyl, -CO<sub>2</sub>C<sub>1-4</sub>alkyl, or -COC<sub>1-4</sub>alkyl;  
 or two R<sup>2</sup> groups are linked together to form a carbocyclic ring that is saturated or  
 30 unsaturated and unsubstituted or substituted by -OH or =O or a heterocyclic ring that is saturated or unsaturated;

- or when P is -CH=CH- and Q is -NR<sup>1</sup>CH<sub>2</sub>-, R<sup>1</sup> and an R<sup>2</sup> are linked together to form a saturated or unsaturated carbocyclic or heterocyclic ring;  
 or when P is -CH=CH- and Q is -NR<sup>1</sup>-, R<sup>1</sup> and an R<sup>2</sup> are linked together to form a saturated or unsaturated carbocyclic or heterocyclic ring, and;
- 5 R<sup>3</sup> is hydrogen, phenylC<sub>1-6</sub> alkyl, C<sub>1-6</sub> alkyl, C<sub>1-6</sub> alkylOCO-, C<sub>1-6</sub>alkylCO-, formyl, CF<sub>3</sub>CO- or C<sub>1-6</sub>alkylSO<sub>2</sub>-, hydroxyC<sub>1-6</sub>alkyl, or C<sub>1-6</sub>alkoxyC<sub>1-6</sub>alkyl.  
 R<sup>7</sup> is hydrogen or C<sub>1-6</sub> alkyl;  
 R<sup>8</sup> is hydrogen or C<sub>1-6</sub> alkyl;  
 R<sup>9</sup> is hydrogen or C<sub>1-6</sub> alkyl;
- 10 R<sup>10</sup> is hydrogen or C<sub>1-6</sub> alkyl;  
 R<sup>11</sup> is hydrogen or C<sub>1-6</sub> alkyl, and;  
 R<sup>12</sup> is hydrogen or C<sub>1-6</sub> alkyl.
2. A compound according to claim 1 wherein
- 15 P is -CH=CH- or Q is CR<sup>1a</sup>=CH and the compound is the E isomer.
3. A compound according to claim 1 or 2 wherein  
 R<sup>1</sup> is hydrogen, fluoro, methyl, ethyl or propyl;  
 R<sup>2</sup> is hydrogen or one or more of methyl, ethyl, *n*-butyl, phenyl, *iso*-propyl,  
 20 *t*-butyl, methoxy, ethoxy, *n*-propoxy, *iso*-propoxy, *n*-butoxy, phenoxy, benzyloxy, bromo, chloro, iodo, fluoro, nitro, cyano, acetyl, pivaloyl, *iso*-butyryl, benzoyl, trifluoromethyl, trifluoromethoxy, trifluoroacetyl, amino, acetylamino, methylthio, oxazolo, methylsulfonyl, *n*-propylsulfonyl, isopropylsulfonyl or dimethylsulfamoyl, and;
- 25 R<sup>3</sup> is hydrogen, methyl, ethyl, propyl, benzyl, *t*-butyloxycarbonyl or trifluoroacetyl.
4. A compound according to any one of claims 1 to 3 wherein  
 R<sup>1</sup> is hydrogen, fluoro or methyl;
- 30 R<sup>2</sup> is hydrogen or one or more of methyl, ethyl, *t*-butyl, methoxy, methoxycarbonyl, methylcarbonyl, ethylcarbonyl, methylamido, acetylamino, methylsulfonyl, oxazole, trifluoromethyl, cyano, chloro, fluoro, or nitro;  
 R<sup>3</sup> is hydrogen, methyl, ethyl, *n*-propyl, benzyl or *t*-butyloxycarbonyl.
- 35 5. A compound of formula (I) according to claim 1 selected from

- E-N-(2-Methyl-1,2,3,4-tetrahydroisoquinolin-7-yl)-3-nitrocinnamide;  
E-N-(2-Methyl-1,2,3,4-tetrahydroisoquinolin-7-yl)-3-trifluoromethylcinnamide;  
E-N-(2-Methyl-1,2,3,4-tetrahydroisoquinolin-7-yl)-3-cinnamide hydrochloride;  
E-N-(2-Methyl-1,2,3,4-tetrahydroisoquinolin-7-yl)-2-methoxycinnamide;  
5 E-N-(2-Methyl-1,2,3,4-tetrahydroisoquinolin-7-yl)-4-chlorocinnamide;  
E-N-(2-Methyl-1,2,3,4-tetrahydroisoquinolin-7-yl)-3-chlorocinnamide;  
E-N-(2-Methyl-1,2,3,4-tetrahydroisoquinolin-7-yl)-3-methoxycinnamide;  
E-N-(2-Methyl-1,2,3,4-tetrahydroisoquinolin-7-yl)- $\alpha$ -methylcinnamide;  
E-N-(2-Methyl-1,2,3,4-tetrahydroisoquinolin-7-yl)-2-chlorocinnamide;  
10 E-N-(2-Methyl-1,2,3,4-tetrahydroisoquinolin-7-yl)-4-methoxycinnamide;  
E-N-(6-methyl-5,6,7,8-tetrahydro[1,6]naphthyridin-3-yl)-3-phenylacrylamide;  
E-3-Furan-2-yl-N-(2-methyl-1,2,3,4-tetrahydroisoquinolin-7-yl)acrylamide;  
E-N-(2-methyl-1,2,3,4-tetrahydroisoquinolin-7-yl)-3-thiophen-2-ylacrylamide;  
E-N-(2-Methyl-1,2,3,4-tetrahydroisoquinolin-7-yl)-2,4-dichlorocinnamide;  
15 Z-N-(2-Methyl-1,2,3,4-tetrahydroisoquinolin-7-yl)-2-methoxycinnamide;  
E-3-Indolin-5-yl-N-(2-Methyl-1,2,3,4-tetrahydroisoquinolin-7-yl)-acrylamide;  
E-3-(1-Methyl-Indolin-2-yl)-N-(2-Methyl-1,2,3,4-tetrahydroisoquinolin-7-yl)-  
acrylamide;  
E-N-(2-Methyl-1,2,3,4-tetrahydroisoquinolin-7-yl)-3-chloro-4-methoxycinnamide;  
20 E-N-(2-Methyl-1,2,3,4-tetrahydroisoquinolin-7-yl)-4-methylsulphonylcinnamide;  
E-N-methyl-3-[2-(2-methyl-1,2,3,4-tetrahydroisoquinolin-7-ylcarbamoyl)vinyl]  
benzamide;  
E-3-(Indazolin-3-yl)-N-(2-Methyl-1,2,3,4-tetrahydroisoquinolin-7-yl)-acrylamide;  
E-N-(2-Methyl-1,2,3,4-tetrahydroisoquinolin-7-yl)-2-methylcinnamide;  
25 E-N-(2-Methyl-1,2,3,4-tetrahydroisoquinolin-7-yl)-2-nitrocinnamide;  
E-N-(2-Methyl-1,2,3,4-tetrahydroisoquinolin-7-yl)-2-trifluoromethylcinnamide;  
E-N-(2-Methyl-1,2,3,4-tetrahydroisoquinolin-7-yl)-2-ethoxycinnamide;  
E-N-(2-Methyl-1,2,3,4-tetrahydroisoquinolin-7-yl)-2-chloro-4-fluorocinnamide;  
E-N-(2-Methyl-1,2,3,4-tetrahydroisoquinolin-7-yl)-2-chloro-6-fluorocinnamide;  
30 E-N-(2-Methyl-1,2,3,4-tetrahydroisoquinolin-5-yl)-4-chlorocinnamide;  
E-N-(2-Methyl-1,2,3,4-tetrahydroisoquinolin-5-yl)-cinnamide;  
E-N-(2-Methyl-1,2,3,4-tetrahydroisoquinolin-5-yl)-3-chlorocinnamide;  
E-N-(2-Methyl-1,2,3,4-tetrahydroisoquinolin-5-yl)-2-chlorocinnamide;  
E-N-(2-Methyl-1,2,3,4-tetrahydroisoquinolin-5-yl)-3-acetylcinnamide;  
35 E-N-(2-Methyl-1,2,3,4-tetrahydroisoquinolin-5-yl)-2-bromocinnamide;  
E-N-(2-Methyl-1,2,3,4-tetrahydroisoquinolin-5-yl)-2-methylcinnamide;  
E-N-(2-Methyl-1,2,3,4-tetrahydroisoquinolin-5-yl)-4-ethoxycinnamide;  
E-N-(2-Methyl-1,2,3,4-tetrahydroisoquinolin-5-yl)-2-methoxycinnamide;  
E-N-(2-Methyl-1,2,3,4-tetrahydroisoquinolin-5-yl)-5-bromo-2-methoxycinnamide;



- E-2-Cyano-N-(2-Methyl-1,2,3,4-tetrahydroisoquinolin-5-yl)cinnamide;  
N-(8-Chloro-2-trifluoroacetyl-1,2,3,4-tetrahydroisoquinolin-7-yl)-2-chlorocinnamide;  
N-(8-Chloro-1,2,3,4-tetrahydroisoquinolin-7-yl)-2-chlorocinnamide;  
5 N-(8-Chloro-2-methyl-1,2,3,4-tetrahydroisoquinolin-7-yl)-2-chlorocinnamide;  
N-(8-Bromo-2-trifluoroacetyl-1,2,3,4-tetrahydroisoquinolin-7-yl)-2-chlorocinnamide;  
E-N-(2-Methyl-1,2,3,4-tetrahydroisoquinolin-7-yl)- $\alpha$ -fluorocinnamide;  
E-N-(8-Bromo-1,2,3,4-tetrahydroisoquinolin-7-yl)-2-chlorocinnamide;  
10 E-N-(8-Bromo-2-methyl-1,2,3,4-tetrahydroisoquinolin-7-yl)-2-chlorocinnamide;  
E-N-(2,4,4-Trimethyl-1,2,3,4-tetrahydroisoquinolin-7-yl)-2-chlorocinnamide;  
E-N-(8-Ethyl-2,4,4-trimethyl-1,2,3,4-tetrahydroisoquinolin-7-yl)-2-methylcinnamide;  
E-N-(8-Ethyl-2,4,4-trimethyl-1,2,3,4-tetrahydroisoquinolin-7-yl)-2-chloro-6-  
15 fluorocinnamide;  
E-N-(8-Ethyl-2,4,4-trimethyl-1,2,3,4-tetrahydroisoquinolin-7-yl)-2-trifluoromethylcinnamide;  
E-N-(8-Ethyl-2,4,4-trimethyl-1,2,3,4-tetrahydroisoquinolin-7-yl)-2-chloro-4-fluorocinnamide;  
20 E-N-(8-Ethyl-2,4,4-trimethyl-1,2,3,4-tetrahydroisoquinolin-7-yl)-2-chlorocinnamide;  
E-N-(1,1,2-Trimethyl-1,2,3,4-tetrahydroisoquinolin-7-yl)-2-chlorocinnamide;  
E-N-(1,2,4,4-Tetramethyl-1,2,3,4-tetrahydroisoquinolin-7-yl)-2-chlorocinnamide;  
E-N-(8-Ethyl-2,4,4-trimethyl-1,2,3,4-tetrahydroisoquinolin-7-yl)-2-  
25 ethoxycinnamide;  
E-N-(8-Chloro-2,4,4-trimethyl-1,2,3,4-tetrahydroisoquinolin-7-yl)-2-chlorocinnamide;  
E-N-(8-Ethyl-2,4,4-trimethyl-1,2,3,4-tetrahydroisoquinolin-7-yl)-3-chloro-4-methoxycinnamide;  
30 E-N-(8-Ethyl-2,4,4-trimethyl-1,2,3,4-tetrahydroisoquinolin-7-yl)-2-cyanocinnamide;  
E-N-(8-Methyl-2,4,4-trimethyl-1,2,3,4-tetrahydroisoquinolin-7-yl)-2-chlorocinnamide;  
E-N-(8-Ethyl-2,4,4-trimethyl-1,2,3,4-tetrahydroisoquinolin-7-yl)-2-  
35 acetylcinnamide;  
E-N-(8-Ethyl-4,4-dimethyl-1,2,3,4-tetrahydroisoquinolin-7-yl)-2-chlorocinnamide;  
E-N-(8-Ethyl-4,4-dimethyl-1,2,3,4-tetrahydroisoquinolin-7-yl)-2-cyanocinnamide;  
E-N-(8-Chloro-2,3,3-trimethyl-1,2,3,4-tetrahydroisoquinolin-7-yl)-2-chlorocinnamide;

E-N-(5-Bromo-1,2,3,4-tetrahydroisoquinolin-7-yl)-2-chlorocinnamide;  
 E-N-(5,6,7,8-Tetrahydro-6-methyl[1,6]naphthyridin-3-yl)-cinnamide, and;  
 E-N-(5,6,7,8-Tetrahydro-6,8,8-trimethyl[1,6]naphthyridin-3-yl)-2-chlorocinnamide.

5

6. A compound of formula (I) according to claim 1 selected from  
 E-N-(4-Methoxyphenyl)-3-(1,2,3,4-tetrahydroisoquinolin-7-yl)acrylamide;  
 E-N-Phenyl-3-(1,2,3,4-tetrahydroisoquinolin-7-yl)acrylamide;

E-N-Phenyl-3-(2-methyl-1,2,3,4-tetrahydroisoquinolin-7-yl)acrylamide;

10 E-N-Phenyl-3-(2-benzyl-1,2,3,4-tetrahydroisoquinolin-7-yl)acrylamide;

E-N-Phenyl-3-(2-n-propyl-1,2,3,4-tetrahydroisoquinolin-7-yl)acrylamide;

E-N-Phenyl-3-(2-ethyl-1,2,3,4-tetrahydroisoquinolin-7-yl)acrylamide;

E-N-(3-Cyanophenyl)-3-(1,2,3,4-tetrahydroisoquinolin-7-yl)acrylamide;

E-N-(3-Cyanophenyl)-3-(2-methyl-1,2,3,4-tetrahydroisoquinolin-7-yl)acrylamide;

15 E-N-(2-Chlorophenyl)-3-(2-methyl-1,2,3,4-tetrahydroisoquinolin-7-yl)acrylamide;

E-N-(2-Methoxyphenyl)-3-(2-methyl-1,2,3,4-tetrahydroisoquinolin-7-yl)acrylamide;

E-N-(3-Methoxyphenyl)-3-(2-methyl-1,2,3,4-tetrahydroisoquinolin-7-yl)acrylamide;

20 E-N-(3-Chlorophenyl)-3-(2-methyl-1,2,3,4-tetrahydroisoquinolin-7-yl)acrylamide;

E-N-(4-Chlorophenyl)-3-(2-methyl-1,2,3,4-tetrahydroisoquinolin-7-yl)acrylamide;

E-N-Methyl-N-benzyl-3-(2-methyl-1,2,3,4-tetrahydroisoquinolin-7-yl)acrylamide;

E-N-(3-Nitrophenyl)-3-(2-methyl-1,2,3,4-tetrahydroisoquinolin-7-yl)acrylamide;

E-N-Methyl-N-phenyl-3-(2-methyl-1,2,3,4-tetrahydroisoquinolin-7-yl)acrylamide;

25 E-N-(3-Carbomethoxyphenyl)-3-(2-methyl-1,2,3,4-tetrahydroisoquinolin-7-yl)acrylamide;

E-N-Methyl-3-[3-(2-methyl-1,2,3,4-tetrahydroisoquinolin-7-yl)acryloylamino]benzamide;

E-N-(3-N-Methylsulphonylphenyl)-3-(2-methyl-1,2,3,4-tetrahydroisoquinolin-7-yl)acrylamide;

30 E-N-(1,3-Oxazol-5-ylphenyl)-3-(2-methyl-1,2,3,4-tetrahydroisoquinolin-7-yl)acrylamide;

E-N-(3-Acetylaminophenyl)-3-(2-methyl-1,2,3,4-tetrahydroisoquinolin-7-yl)acrylamide;

35 E-N-(3-Ethylphenyl)-3-(2-methyl-1,2,3,4-tetrahydroisoquinolin-7-yl)acrylamide;


E-N-(3-Methylphenyl)-3-(2-methyl-1,2,3,4-tetrahydroisoquinolin-7-yl)acrylamide;

E-N-(3-tert-Butylphenyl)-3-(2-methyl-1,2,3,4-tetrahydroisoquinolin-7-yl)acrylamide;


E-N-(4-Fluorophenyl)-3-(2-methyl-1,2,3,4-tetrahydroisoquinolin-7-yl)acrylamide;

- E-N-(4-Methoxyphenyl)-3-(2-methyl-1,2,3,4-tetrahydroisoquinolin-7-yl)acrylamide;  
 E-N-(4-Carbomethoxyphenyl)-3-(2-methyl-1,2,3,4-tetrahydroisoquinolin-7-yl)acrylamide;  
 5 E-N-(4-Cyanophenyl)-3-(2-methyl-1,2,3,4-tetrahydroisoquinolin-7-yl)acrylamide;  
 E-N-(4-Nitrophenyl)-3-(2-methyl-1,2,3,4-tetrahydroisoquinolin-7-yl)acrylamide;  
 E-N-(4-Methylphenyl)-3-(2-methyl-1,2,3,4-tetrahydroisoquinolin-7-yl)acrylamide;  
 E-N-(3-Methoxy-5-trifluoromethylphenyl)-3-(2-methyl-1,2,3,4-tetrahydroisoquinolin-7-yl)acrylamide;  
 10 E-1-(3,4-Dihydro-1H-isoquinolin-2-yl)-3-(2-methyl-1,2,3,4-tetrahydroisoquinolin-7-yl)propenone;  
 E-1-(3,4-Dihydro-2H-quinolin-1-yl)-3-(2-methyl-1,2,3,4-tetrahydroisoquinolin-7-yl)propenone;  
 E-1-(3,3-Dimethyl-2,3-dihydroindol-1-yl)-3-(2-methyl-1,2,3,4-tetrahydroisoquinolin-7-yl)propenone, and;  
 15 E-1-(2,3-Dihydroindol-1-yl)-3-(2-methyl-1,2,3,4-tetrahydroisoquinolin-7-yl)propenone.


7. A process for the preparation of compounds of formula (I), which  
 20 comprises  
 (a) for compounds of formula (I) in which P is -NH- and Q is -CR<sup>1</sup>=CH-, reacting a compound of formula (II)




- 25 with a compound of formula (III)



- 30 or,  
 (b) for compounds of formula (I) in which P is -CH=CH- and Q is -NR<sup>1</sup>-, reacting a compound of formula (IV)



with a compound of formula (V)




where  $R^{1A}$ ,  $R^{2A}$ ,  $R^{3A}$ ,  $R^{7A}$ ,  $R^{8A}$ ,  $R^{9A}$ , and  $R^{10A}$  are independently  $R^1$ ,  $R^2$ ,  $R^3$ ,  $R^7$ ,  $R^8$ ,  $R^9$ , and  $R^{10}$  as defined for formula (I) or a group or groups convertible thereto; Z, X and Y are as defined for formula (I); and L is OH or a halogen;

and where required converting an  $R^{1A}$ ,  $R^{2A}$ ,  $R^{3A}$ ,  $R^{7A}$ ,  $R^{8A}$ ,  $R^{9A}$ , or  $R^{10A}$  group to an  $R^1$ ,  $R^2$ ,  $R^3$ ,  $R^7$ ,  $R^8$ ,  $R^9$ , or  $R^{10}$  group;

converting one  $R^1$ ,  $R^2$ ,  $R^3$ ,  $R^7$ ,  $R^8$ ,  $R^9$ , or  $R^{10}$  group to another  $R^1$ ,  $R^2$ ,  $R^3$ ,  $R^7$ ,  $R^8$ ,  $R^9$ , or  $R^{10}$  group;

converting a salt product to the free base or another pharmaceutically acceptable salt, or converting a free base product to a pharmaceutically acceptable salt.

8. A compound of formula (XII)



wherein  $R^{3A}$  is  $R^3$  as defined in claim 1 or a group convertible thereto and M is a leaving group such as halogen, especially iodo, or tosylate.

9. A pharmaceutical composition for use in the treatment and/or prevention of anxiety, mania, depression, panic disorders and/or aggression, disorders associated with a subarachnoid haemorrhage or neural shock, the effects

associated with withdrawal from substances of abuse such as cocaine, nicotine, alcohol and benzodiazepines, disorders treatable and/or preventable with anti-convulsive agents, such as epilepsy including post-traumatic epilepsy, Parkinson's disease, psychosis, migraine, cerebral ischaemia, Alzheimer's disease and other  
5 degenerative diseases such as Huntingdon's chorea, schizophrenia, obsessive compulsive disorders (OCD), neurological deficits associated with AIDS, sleep disorders (including circadian rhythm disorders, insomnia & narcolepsy), tics (e.g. Gilles de la Tourette's syndrome), traumatic brain injury, tinnitus, neuralgia, especially trigeminal neuralgia, neuropathic pain, dental pain, cancer pain,  
10 inappropriate neuronal activity resulting in neurodysthesias in diseases such as diabetes, multiple sclerosis (MS) and motor neurone disease, ataxias, muscular rigidity (spasticity), temporomandibular joint dysfunction, and amyotrophic lateral sclerosis (ALS) which comprises a compound of formula (I) as defined in claim 1, or a pharmaceutically acceptable salt or solvate thereof, and a pharmaceutically  
15 acceptable carrier.

10. A method of treatment and/or prevention of anxiety, mania, depression, panic disorders and/or aggression, disorders associated with a subarachnoid haemorrhage or neural shock, the effects associated with withdrawal from  
20 substances of abuse such as cocaine, nicotine, alcohol and benzodiazepines, disorders treatable and/or preventable with anti-convulsive agents, such as epilepsy including post-traumatic epilepsy, Parkinson's disease, psychosis, migraine, cerebral ischaemia, Alzheimer's disease and other degenerative diseases such as Huntingdon's chorea, schizophrenia, obsessive compulsive disorders  
25 (OCD), neurological deficits associated with AIDS, sleep disorders (including circadian rhythm disorders, insomnia & narcolepsy), tics (e.g. Gilles de la Tourette's syndrome), traumatic brain injury, tinnitus, neuralgia, especially trigeminal neuralgia, neuropathic pain, dental pain, cancer pain, inappropriate neuronal activity resulting in neurodysthesias in diseases such as diabetes,  
30 multiple sclerosis (MS) and motor neurone disease, ataxias, muscular rigidity (spasticity), temporomandibular joint dysfunction, and amyotrophic lateral sclerosis (ALS) comprising administering to the sufferer in need thereof an effective or prophylactic amount of a compound of formula (I) as defined in claim 1, or a pharmaceutically acceptable salt or solvate thereof.

35  
11. Use of a compound of formula (I) as defined in claim 1, or a pharmaceutically acceptable salt or solvate thereof, for the manufacture of a medicament for the treatment and/or prevention of anxiety, mania, depression, panic disorders and/or aggression, disorders associated with a subarachnoid

- haemorrhage or neural shock, the effects associated with withdrawal from substances of abuse such as cocaine, nicotine, alcohol and benzodiazepines, disorders treatable and/or preventable with anti-convulsive agents, such as epilepsy including post-traumatic epilepsy, Parkinson's disease, psychosis, migraine, cerebral ischaemia, Alzheimer's disease and other degenerative diseases
- 5 such as Huntingdon's chorea, schizophrenia, obsessive compulsive disorders (OCD), neurological deficits associated with AIDS, sleep disorders (including circadian rhythm disorders, insomnia & narcolepsy), tics (e.g. Gilles de la
- 10 Tourette's syndrome), traumatic brain injury, tinnitus, neuralgia, especially trigeminal neuralgia, neuropathic pain, dental pain, cancer pain, inappropriate neuronal activity resulting in neurodysthesias in diseases such as diabetes, multiple sclerosis (MS) and motor neurone disease, ataxias, muscular rigidity (spasticity), temporomandibular joint dysfunction, and amyotrophic lateral sclerosis (ALS).
- 15
12. Use of a compound of formula (I) as defined in claim 1, or a pharmaceutically acceptable salt or solvate, thereof as a therapeutic agent, in particular for the treatment and/or prevention of anxiety, mania, depression, panic disorders and/or aggression, disorders associated with a subarachnoid
- 20 haemorrhage or neural shock, the effects associated with withdrawal from substances of abuse such as cocaine, nicotine, alcohol and benzodiazepines, disorders treatable and/or preventable with anti-convulsive agents, such as epilepsy including post-traumatic epilepsy, Parkinson's disease, psychosis, migraine, cerebral ischaemia, Alzheimer's disease and other degenerative diseases
- 25 such as Huntingdon's chorea, schizophrenia, obsessive compulsive disorders (OCD), neurological deficits associated with AIDS, sleep disorders (including circadian rhythm disorders, insomnia & narcolepsy), tics (e.g. Gilles de la Tourette's syndrome), traumatic brain injury, tinnitus, neuralgia, especially trigeminal neuralgia, neuropathic pain, dental pain, cancer pain, inappropriate
- 30 neuronal activity resulting in neurodysthesias in diseases such as diabetes, multiple sclerosis (MS) and motor neurone disease, ataxias, muscular rigidity (spasticity), temporomandibular joint dysfunction, and amyotrophic lateral sclerosis (ALS).

## INTERNATIONAL SEARCH REPORT

International Application No

PC1/EP 99/05583

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07D217/04 C07D471/04 A61K31/472 A61K31/4725 A61K31/4375  
 C07D405/12 C07D409/12 C07D401/12 C07D217/06 C07D217/02  
 C07D413/12 C07D401/06 //(C07D471/04,221:00,221:00)

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07D A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	MATHISON I W ET AL: "SYNTHESIS AND HYPOTENSIVE PROPERTIES OF TETRAHYDROISOQUINOLINES" JOURNAL OF MEDICINAL CHEMISTRY, US, AMERICAN CHEMICAL SOCIETY. WASHINGTON, vol. 16, no. 4, page 332-336 XP002040786 ISSN: 0022-2623 see page 332, scheme 1 and page 334 experimental section the whole document	8
A	---	1,9
A	GB 1 164 192 A (FARBWERKE HOECHST AG) 17 September 1969 (1969-09-17) page 1, line 6 - line 8; claims 2,17	
A	WO 97 48683 A (SMITHKLINE BEECHAM PLC) 24 December 1997 (1997-12-24) claims	1,9-12

☐ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

## \* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"Z" document member of the same patent family

Date of the actual completion of the international search

12 November 1999

Date of mailing of the international search report

29/11/1999

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
 NL - 2280 HV Rijswijk  
 Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
 Fax: (+31-70) 340-3016

Authorized officer

Henry, J

# INTERNATIONAL SEARCH REPORT

I. International application No.

PCT/EP 99/05583

## Box I Observations where certain claims were found unsearchable (Continuation of Item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.: 10  
because they relate to subject matter not required to be searched by this Authority, namely:  
Remark: Although claim 10  
is directed to a method of treatment of the human/animal  
body, the search has been carried out and based on the alleged  
effects of the compound/composition.
2. ☐ Claims Nos.:  
because they relate to parts of the International Application that do not comply with the prescribed requirements to such  
an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

## Box II Observations where unity of invention is lacking (Continuation of Item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all  
searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment  
of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report  
covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is  
restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

☐ The additional search fees were accompanied by the applicant's protest.

☐ No protest accompanied the payment of additional search fees.



# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/EP 99/05583

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
GB 1164192 A	17-09-1969	DE 1670694 A	03-12-1970
		DE 1670848 A	08-04-1971
		DE 1670849 A	25-02-1971
		AT 281827 B	10-06-1970
		AT 289804 B	15-03-1971
		BE 698033 A	06-11-1967
		CH 487893 A	31-03-1970
		CH 501628 A	15-01-1971
		CH 487892 A	31-03-1971
		DK 123598 B	10-07-1972
		FR 6496 M	25-11-1968
		FR 6646 M	20-01-1969
		FR 1524487 A	02-09-1968
		NL 6706322 A,B	06-11-1967
		SE 330170 B	09-11-1970
		US 3577424 A	04-05-1971
		AT 282634 B	10-07-1970
WO 9748683 A	24-12-1997	AU 3259597 A	07-01-1998
		CA 2258238 A	24-12-1997
		CZ 9804164 A	16-06-1999
		EP 0906283 A	07-04-1999
		NO 985891 A	16-12-1998
		NZ 332757 A	29-06-1999
		PL 330465 A	24-05-1999